arXiv:1101.1022v2 [math.CO] 16 Jul 2012

ON CHIROTOPES OF FINITE PLANAR FAMILIES OF PAIRWISE DISJOINT CONVEX BODIES

LUC HABERT AND MICHEL POCCHIOLA

Abstract. We extend the classical characterization of chirotopes of finite planar families of points—which says that a map χ defined on the set of triples of a finite indexing set I is the chirotope of a finite planar family of points if and only if for every 3-, 4-, and 5-subset J of I the restriction of χ to the set of triples of J is the chirotope of a finite planar family of points—to chirotopes of finite planar families of pairwise disjoint convex bodies. Our main tool is the polarity map, i.e., the map that assigns to a convex body the set of lines missing its interior, from which we derive the key notion of arrangements of double pseudolines, introduced for the first time in this paper.

Keywords. Convexity, discrete geometry, projective planes, pseudoline arrangements, chirotopes.

Abbreviated version in Proc. 25th Annu. ACM Sympos. Comput. Geom. (SCG09), pages 314–323, June 2009, Aahrus, Denmark. A preliminary abbreviated version of that paper appears in the Abstracts 12th European Workshop Comput. Geom. pages 211–214, 2006, Delphes, and a poster version was presented at the Workshop on Geometric and Topological Combinatorics (satellite conference of ICM 2006), September 2006, Alcala de Henares, Spain.

Date: July 17, 2012.
MP was partially supported by the TEOMATRO grant ANR-10-BLAN 0207.

Contents

1. Introduction 3
1.1. Cross surfaces and projective planes 3
1.2. Definitions and main results 6
1.3. Organization of the paper 11
2. Homotopy theorem 11
2.1. Thin arrangements of double pseudolines 11
2.2. The pumping lemma 12
2.3. Enumeration of arrangements 16
3. Geometric representation theorem 18
3.1. Nodes and cycles of an arrangement 18
3.2. Factorization of the duality transform, raiponces 21
3.3. Stability by mutations 22
4. Cycles, cocycles and chirotopes 29
4.1. Side cycles 29
4.2. Chirotopes 36
4.3. Cocycles 37
5. Axiomatic characterization 45
5.1. Arrangements of genus $1, 2, \ldots$ 46
5.2. Pumping lemma and mutations 56
6. An extension and a refinement 61
7. Conclusion and open problems 65
Acknowledgments. 67
References 68
Appendix A. Arrangements of pseudolines 70
Appendix B. Chirotopes of finite planar families of points 75
Appendix C. Convexity in neutral and projective planes 78
C.1. Background material on neutral and affine planes 78
C.2. Convexity and duality in (real two-dimensional) neutral planes 79
C.3. Convexity and duality in projective planes 83

1. INTRODUCTION

The term planar in the title makes reference to real two-dimensional projective planes. We review what we need of the basics of real two-dimensional projective planes and especially the notion of convex body before introducing the notion of chirotope, explaining the main result of the paper and the main lines of its proof.

1.1. **Cross surfaces and projective planes.** We assume the reader familiar with the basic notions of algebraic or combinatorial topology like homeomorphism, homotopy, fundamental group, covering, etc., that can be found for example in [30, chap. 0 and 1]. The following standard notions, basic results and terminology associated with projective planes will be used throughout the paper; they are mainly taken from [40, 24, 36, 39]

(1) A *closed (open) topological disk* or *closed (open) two-cell* is a topological space homeomorphic to the unit closed (open) disk of $\mathbb{R}^2$; an *orientation* of a topological disk is a one-to-one parametrization of the topological disk by the unit disk of $\mathbb{R}^2$, defined up to direct homeomorphism; and an *oriented topological disk* is a topological disk endowed with an orientation; orientations will be indicated in our drawings by a little oriented circle in the interior of the disk or by an arrow on its boundary.

(2) A *cross surface*[1] is a topological space homeomorphic to the "standard" cross surface $\mathbb{RP}^2$, quotient of the unit sphere $\mathbb{S}^2$ of $\mathbb{R}^3$ under identification of antipodal points; cross surfaces will be represented in our drawings by circular diagrams with antipodal boundary points identified.

(3) An *open crosscap* or *open Möbius strip* is a topological space homeomorphic to a cross surface with one point or one closed topological disk deleted; an open crosscap is a noncompact surface and its one-point compactification (the space obtained by adding to the crosscap a point at infinity) is a cross surface.

(4) A *pseudocircle* is a simple closed curve embedded in a cross surface; the connected components of the complement of a pseudocircle in its underlying cross surface are called its *open sides*, or simply its *sides*. An *oriented pseudocircle* is a pseudocircle endowed with an *orientation* (i.e., a one-to-one parametrization of the pseudocircle by $\mathbb{S}^1$, defined up to direct homeomorphism), indicated in our drawings by an arrow; as usual the intersection of two oriented pseudocircles is the intersection of their unoriented versions.

(5) A *pseudoline* is a non-separating pseudocircle and a *double pseudoline* or *pseudo-oval* is a separating pseudocircle. There is a unique isomorphism class of pseudolines, i.e., given two pseudolines, one is the image of the other by a homeomorphism of their underlying cross surfaces; in particular the complement of a pseudoline is an open two-cell. Similarly for double pseudolines: there is a unique isomorphism class of double pseudolines and the complement of a double pseudoline has two connected components : an open two-cell and an open crosscap; the *core* pseudolines of a double pseudoline are the pseudolines contained in its crosscap side. Figure 1 depicts a cross surface represented by a circular diagram with antipodal points identified, a pseudoline, a double pseudoline with one of its

---

[1] We follow the J. H. Conway' proposition to call a sphere with one crosscap a *cross surface*; cf. [21].

core pseudolines, and an oriented double pseudoline with one of its core oriented pseudolines.



FIGURE 1. A cross surface represented by a circular diagram with antipodal points identified, a pseudoline, a double pseudoline with one of its core pseudolines and with its disk side in white, and an oriented double pseudoline with one of its core oriented pseudolines.

(6) A *projective plane* is a topological point-line incidence geometry $(\mathcal{P}, \mathcal{L})$ whose point space $\mathcal{P}$ is a cross surface, whose line space $\mathcal{L}$ is a subspace of the space of pseudolines of $\mathcal{P}$, and whose incidence relations are the membership relations; as usual the dual of a point p of a projective plane is denoted p^* and is defined as its set of incident lines. The *duality principle* for projective planes asserts that the dual $(\mathcal{L}, \mathcal{P}^*)$ of a projective plane $(\mathcal{P}, \mathcal{L})$ is still a projective plane, i.e., $\mathcal{L}$ is a cross surface and $\mathcal{P}^*$, the set of p^* as p ranges over $\mathcal{P}$, is a subspace of the space of pseudolines of $\mathcal{L}$; in particular the dual of a finite set of points is an *arrangement of pseudolines*, i.e., a finite set of pseudolines (living in the same cross surface) that intersect pairwise in exactly one point; the basics of pseudoline arrangements used in the paper are reviewed in Appendix A. A projective plane is isomorphic to its bidual via the map that assigns to a point its dual and to a line the set of duals of its points.

(7) The standard projective plane is defined as the standard cross surface $\mathbb{RP}^2$ together with the image of the space of great circles of $\mathbb{S}^2$ under the canonical projection $\mathbb{S}^2 \to \mathbb{RP}^2$. (Equivalently the standard projective plane can be defined as the projective completion of the Euclidean plane.) The standard projective plane is isomorphic to its dual via the map φ that assigns to the point $(u, v, w) \in \mathbb{S}^2$ the great circle with equation $ux + vy + wz = 0$ and that assigns to the great circle with equation $ux+vy+wz = 0$,$(u, v, w) \in \mathbb{S}^2$, the pencil of circles through the point (u, v, w).

A *convex body* is a closed topological disk embedded in the point space of a projective plane whose intersection with any line of the plane is a (necessarily closed) line-segment; the *polar* of a convex body U, denoted $U^\diamond$, is the set of lines of the plane missing the interior of the convex body and its *dual*, denoted U^*, is the set of lines of the plane intersecting the body but not its interior, *tangents* for short. For example, for $(u, v, w) \in \mathbb{S}^2$ and $h \in (0, 1)$, the disk with equation

$$|ux + vy + wz| \geq (1 - h^2)^{1/2} \tag{1}$$

of the standard projective plane is a convex body, its polar is the disk with equation $|ux+vy+wz| \geq h$, and its dual is the circle with equation $|ux+vy+wz| = h$. Similarly for finitely generated (pointed and full-dimensional) cones of the standard projective plane: the polar of the cone generated by the vectors $(u_i, v_i, w_i) \in \mathbb{S}^2$, $w_i > 0$, is the polyhedral cone intersection of the half-spaces $u_i x + v_i y + w_i z \geq 0$, $z \geq 0$. As illustrated in these examples the polar of a convex body is a convex body of the dual projective plane and its dual is the boundary of its polar, hence a double pseudoline; furthermore polarity extends to oriented convex bodies : the polar of an oriented convex body has a natural orientation, inherited from the orientation of the body, compatible with the reorientation operation (in the case where there is exactly one tangent through each boundary point and only one touching point per tangent the orientation of the polar inherited from the orientation f is simply defined as the extension to the unit closed disk of the map that assigns to $u \in \mathbb{S}^1$ the tangent to the convex body through the boundary point $f(u)$ of U; the general case follows once observed that the set of boundary points through which pass a proper interval of tangents and the set of proper line-segments included in the boundary are both countable); as far as we know no proof of these basic facts are available in the literature [13, 31, 7, 8, 9, 15, 14, 12, 11, 33]; for completeness we offer proofs in Appendix C. The last basic result to be mentioned is a complete characterization, up to homeomorphism, of the dual arrangement of a pair of disjoint convex bodies of a projective plane.

Theorem 1. *Up to homeomorphism the dual arrangement of a pair of disjoint convex bodies of a projective plane is the unique arrangement of two double pseudolines that intersect in four transversal intersection points and induce a cellular decomposition of their underlying cross surface; in particular two disjoint convex bodies share exactly four common tangents, the arrangement of these four tangents is simple, and the set of lines missing the two bodies is nonempty.* □

Proof. See Appendix C. □

Figure 2 depicts an indexed family of two disjoint oriented convex bodies and its dual arrangement; in this figure each body is marked with an interior point; modulo the choice of the nature of the contacts (points or line-segments) between the bodies and the tangents the left diagram is unique up to homeomorphism; the right diagram is



FIGURE 2. An indexed family of two disjoint oriented and marked convex bodies and its dual arrangement.

unique up to homeomorphism, the automorphism group of its cell poset is trivial, $\mathbb{Z}_2$ or the dihedral group of order 8 depending on whether orientation and indexation of the pseudocircles are both taken into account, the orientation of the pseudocircles is taken into account but not their indexation, or neither the orientation nor the indexation are taken into account; observe that this diagram does not encode the nature of the contacts between the convex bodies and their common tangents. The four common tangents of two disjoint convex bodies will be called their *bitangents*.

1.2. **Definitions and main results.** Throughout the paper we use the words *configuration of convex bodies* for a finite family of pairwise disjoint convex bodies of a projective plane and we use the words *arrangement of double pseudolines* for a finite family of double pseudolines of a cross surface with the property that its subfamilies of size two are homeomorphic to the dual arrangement of a (hence any) configuration of two convex bodies; cf. Theorem 1. We extend in the natural way the basic terminology of arrangements of pseudolines to arrangements of double pseudolines: in particular

(1) the *isomorphism class* of an arrangement is its set of homeomorphic images : in other terms, two arrangements are called *isomorphic* if one is the image of the other by a homeomorphism of their underlying cross surfaces;
(2) the *face poset* of an arrangement is its set of closed cells ordered by containment; the *flags* of an arrangement are the maximal chains (or maximal simplices of the first barycentric subdivision) of its face poset and its 0-, 1- and 2-*flag operators* are the involutive operators, defined on the set of flags, that exchange two flags that differ only by their 0-, 1-, and 2-cells, respectively; arrangements of double pseudolines are conveniently represented by their face posets or, equivalently, by their flag operators, in view of the two following properties:
    (a) two arrangements are isomorphic if and only if their face posets are isomorphic, cf. [2, Appendix 4.7]; and
    (b) the group of automorphisms of an arrangement (by definition quotient of the group of self-homeomorphisms of the arrangement by its subgroup of self-homeomorphisms isotopic to the identity map) is isomorphic to the group of automorphisms of its face poset or, equivalently, to the centralizer of the flag operators in the group of permutations of the flags;
(3) the *chirotope* of an indexed arrangement of oriented double pseudolines is the map that assigns to each triple of the indexing set of the arrangement the isomorphism class of the subarrangement indexed by this triple;
(4) a *mutation* is a homotopy of arrangements during which only one of the curves of the arrangement is moving and only one of the incidence relations between the moving curve and the vertices of the cell complex induced by the other curves changes its value, swapping from false to true (first case) or from true to false (second case): In the first case we speak of a *merging* mutation and in second case we speak of a *splitting* mutation.

Example 1. Figure 3 shows four examples of arrangements of three double pseudolines; the first example is isomorphic to the family of 3 circles of the standard projective plane with equations $|x| = |y| = |z| = r$, $0 < r < 1/\sqrt{3}$, and its face poset is that of the hemi-rhombicubeoctahedron (the projective version of the rhombicubeoctahedron, one of the

13 Archimedean solids); the second example is an indexed and oriented version of the first



FIGURE 3. Examples of arrangements of double pseudolines.

example; the third example is isomorphic to the family of 3 circles $|x| = |y| = |z| = 1/\sqrt{3}$ of the standard projective plane and its face poset is obtained from that of the hemi-cube (the projective version of the cube) by replacing each of its 1-cells by a digon; the fourth example is derived from the third one by a splitting mutation and, conversely, the third one is derived from the fourth one by a merging mutation.

Example 2. Figure 4 depicts the face poset of an arrangement of two double pseudolines (top row) and the face poset of the third example of Figure 3 (bottom row) under the guise of their first barycentric subdivisions (left row) and under the guise of their *flag diagrams* (right row) : the flag diagram of the face poset of an arrangement is the 3-valent graph whose nodes are the flags of the cell poset (maximal chain of the cell poset or, equivalently, maximal simplices of its first barycentric subdivision) and whose edges are the pairs of adjacent flags, each edge being labeled by the numeral $0, 1$ or 2 depending on whether the adjacent flags differ by a 0, 1-, or 2-cell. (One can also think a flag diagram as the Cayley graph of the group generated by the flag operators, denoted σ_0, σ_1 and σ_2 in the sequel.) Each diagram is labeled at its left bottom corner by a symbol to named it and at its right bottom corner by the size of its automorphism group. The automorphism group of the arrangement A is a dihedral group of order 8, group of automorphisms of the square; two generators τ_1 and τ_2 of which $\tau_1^2 = \tau_2^2 = 1$ and $(\tau_2\tau_1)^4 = 1$ is a complete set of relations are, for example, those defined by $\tau_1(F) = \sigma_1(F)$ and $\tau_2(F) = \sigma_0(F)$ (since the face poset of an arrangement is flag-connected an automorphism is defined by the image of one flag) where F is any one of the 8 flags of the tetragon intersection of the crosscap sides of the double pseudolines of the arrangement (they are labeled $1, 2, \ldots, 8$ in the figure). The automorphism group of the arrangement Z is S_4, the group of automorphisms of the hemicube; three generators τ_1, τ_2, τ_3 of which $\tau_1^2 = \tau_2^2 = \tau_3^3 = 1$, $(\tau_1\tau_2)^4 = 1$, $\tau_1\tau_3 = \tau_3^2\tau_1$ and $\tau_2\tau_3 = \tau_3(\tau_1\tau_2)^2$ is a complete set of relations are, for example, those defined by $\tau_1(F) = \sigma_1(F), \tau_2(F) = \sigma_0(F)$, and $\tau_3(F) = \sigma_1\sigma_2\sigma_1\sigma_2(F)$ where F is any one of the 3×8 flags of the 3 tetragons of the arrangement (each tetragon is the intersection of the crosscap sides of a pair of double pseudolines; the 24 flags are labeled $1, 1', 1'', \ldots, 8, 8', 8''$ in the figure).

Our definition of chirotopes of configurations of convex bodies is a natural extension of the classical definition of chirotopes of configurations of points of the standard projective plane; cf. Appendix B. Let Δ be an indexed configuration of oriented convex bodies of a projective plane $(\mathcal{P}, \mathcal{L})$, and let τ be a line of $(\mathcal{P}, \mathcal{L})$. We denote by $\omega_\tau : \mathcal{P} \to \mathcal{P}/\mathcal{R}_\tau$



Figure 4. The face poset of an arrangement

the quotient map relative to the equivalence relation $\mathcal{R}_\tau$ whose equivalence classes not reduced to a point are the proper line segments of $\Delta \cap \tau$ and we define

1. the *cocycle of* Δ *at* τ or the *cocycle of the pair* (Δ, τ) as the homeomorphism class of the image of the pair (Δ, τ) under ω_τ, i.e., the set of $(\varphi\omega_\tau\Delta, \varphi\omega_\tau\tau)$ as φ ranges over the set of homeomorphisms with domain $\mathcal{P}/\mathcal{R}_\tau$ (ω_τ is there to make no difference between a tangent touching a convex body along a line segment or along a point);
2. a *bitangent cocycle* or *zero-cocycle* as a cocycle at a bitangent;
3. the *isomorphism class of* Δ as the set of configurations Δ' that have the same set of bitangent cocycles as Δ, hence the same set of cocycles as Δ (use a simple perturbation argument); and
4. the *chirotope of* Δ as the map that assigns to each triple J of the indexing set of Δ the isomorphism class of the subfamily indexed by J.

Figure 5 depicts, up to reorientation and reindexation of the bodies, the bitangent cocycles of configurations of two and three convex bodies : each circular diagram is labeled at its right bottom corner with its number of reoriented and reindexed versions, and at its left bottom corner with its "natural" coding (a set of words on the alphabet obtained by adjoining to the indexing set of the configuration the negatives of its elements and the extra symbol •, with the convention that the negative of an index α is denoted $\overline{\alpha}$ instead of $-\alpha$) that we will define explicitly in the body text. One can read from these data that the number of bitangent cocycles of indexed configurations of two oriented convex bodies on a given set of two indices is exactly the number (4) of bitangents of a pair of disjoint convex bodies; and that the number of bitangent cocycles of indexed configurations of three oriented convex bodies on a given set of three indices is 104.


12•• 4
1•2•3• 8
123••• 24
$\overline{3}$123•• 24
1$\overline{3}$2•3• 24
12••, 3 24

FIGURE 5. The bitangent cocycles of configurations of convex bodies indexed by $1, 2$ and $1, 2, 3$, respectively.

Figure 6 depicts an indexed configuration of three oriented convex bodies together with its bitangents (the bitangents are all tritangents and are labeled A, B, C and D), its bitangent cocycles, and its dual arrangement.

The four main results of the paper, the first one being instrumental to prove the other three, are the following.

Theorem 2. *Any two arrangements of double pseudolines of the same size and living in the same cross surface are homotopic via a finite sequence of mutations followed by an isotopy; in other terms, mutation graphs are connected.* □

Theorem 3. *The map that assigns to an indexed configuration of oriented convex bodies the isomorphism class of its dual arrangement is compatible with the isomorphism relation on indexed configurations of oriented convex bodies and the induced quotient map is one-to-one and onto, i.e., any arrangement of double pseudolines is isomorphic to the dual arrangement of a configuration of convex bodies.* □


A
B
C
D
1
2
3
$1\bullet\overline{2}\bullet\overline{3}\bullet$
$1\bullet\overline{3}\bullet 2\bullet$
$1\bullet 2\bullet 3\bullet$
$1\bullet 3\bullet\overline{2}\bullet$

FIGURE 6. An indexed configuration of three oriented convex bodies, its bitangent cocycles, and its dual arrangement.

For example, Theorem 3 asserts that the set of bitangent cocycles $\{1\bullet\overline{2}\bullet\overline{3}\bullet\}$, $\{1\bullet\overline{3}\bullet 2\bullet\}$, $\{1\bullet 2\bullet 3\bullet\}$, $\{1\bullet 3\bullet\overline{2}\bullet\}$ of the configuration of three convex bodies depicted in the left part of Figure 6 is a coding of the isomorphism class of its dual arrangement.

Theorem 4. *The map that assigns to an isomorphism class of indexed arrangements of oriented double pseudolines its chirotope is one-to-one and its range is the set of map χ defined on the set of triples of a finite set I such that for every 3-, 4-, and 5-subset J of I the restriction of χ to the set of triples of J is the chirotope of an indexed arrangement of oriented double pseudolines.* ☐

Theorem 5. *The map that assigns to an isomorphism class of finite indexed configuration of oriented convex bodies its chirotope is one-to-one and its range is the set of map χ defined on the set of triples of a finite set I such that for every 3-, 4-, and 5-subset J of I the restriction of χ to the set of triples of J is the chirotope of a finite indexed configuration of oriented convex bodies.* ☐

Note that Theorem 5 is a simple combination of Theorems 3 and 4. The well-informed reader will observe that the four above theorems are direct extensions of classical theorems on arrangements of pseudolines and/or configurations of points : Theorem 2 is a direct extension of the Ringel homotopy theorem for arrangements of pseudolines [37]; the onto part of Theorem 3 is a direct extension of a reformulation, using the duality principle for projective planes as explained in Appendix A, of the enlargement theorem for pseudoline arrangements of Goodman, Pollack, Wenger and Zamfirescu [24]; Theorem 4 is a direct extension of the three-dimensional case of the Folkman and Lawrence axiomatic characterization of the class of isomorphism classes of arrangements of pseudohyperplanes [20]; and Theorem 5 is a direct extension of the classical characterization of chirotopes of finite planar families of point mentioned in the abstract; cf. Appendix B.

1.3. **Organization of the paper.** The paper is organized as follows. In Section 2 we prove that mutation graphs are connected; we give the full list of isomorphism classes of simple arrangements of three double pseudolines; and we extract from this list the set of *martagons* on three and four double pseudolines: martagons are arrangements that show up naturally in the proof that the isomorphism class of an indexed arrangement of oriented double pseudolines depends only on its chirotope; cf. first part of Theorem 4. In Section 3 we show, using the connectedness of mutation graphs proved in Section 2, that any arrangement of double pseudolines is isomorphic to the dual arrangement of a configuration of convex bodies. In section 4 we prove that the duality map is compatible with the isomorphism relations on the set of configurations of convex bodies and on the set of arrangements of double pseudolines, and that the induced quotient map is one-to-one. In Section 5 we prove the axiomatic characterization of chirotopes of double pseudoline arrangements : the proof is based on an extension of the notion of double pseudoline arrangements obtained by relaxing the condition on the arrangement which says that the genus of its underlying (nonorientable) surface is 1. We also provide a comprehensive description of the class of arrangements of 5 double pseudolines in terms of the transitivity of certain relations defined by the arrangements of 4 double pseudolines, reminiscent of the 3-terms Grassmann-Plücker relations for arrangements of pseudolines. In Section 6 we discuss *arrangements of pseudocircles* (as natural extensions of both arrangements of pseudolines and arrangements of double pseudolines), *crosscap or Möbius arrangements* and their *fibrations* (as dual arrangements of affine configurations of convex bodies with, in particular, a positive answer to a question of Goodman and Pollack about the realizability of their double permutation sequences by affine configurations of pairwise disjoint convex bodies). We conclude in the sixth and last section with a list of open problems suggested by this research.

2. HOMOTOPY THEOREM

In this section we prove that any two arrangements of double pseudolines with the same number of double pseudolines and living in the same cross surface are homotopic via a finite sequence of mutations followed by an isotopy. We proceed into two steps

1. firstly, in order to benefit from the Ringel's homotopy theorem for arrangements of pseudolines, we embed the class of isomorphism classes of simple arrangements of pseudolines into the class of isomorphism classes of arrangements of double pseudolines; the embedding is canonical and is based on the notion of *thin* arrangement of double pseudolines;
2. secondly (and this is the core of our proof) we introduce a 'pumping' device to come down to the case of arrangements of pseudolines.

We also provide representatives of the isomorphism classes of simple arrangements of three double pseudolines and we use these representatives to compute the full list of *martagons* on three and four double pseudolines: martagons are arrangements that play a special rôle in the proof that the isomorphism class of an indexed arrangement of oriented double pseudolines depends only on its chirotope.

2.1. **Thin arrangements of double pseudolines.** A simple arrangement of double-pseudolines is termed *thin* if the crosscap sides of its double pseudolines are free of vertices, and a thin arrangement of double pseudolines Γ^* is termed a *double* of a simple arrangement of pseudolines Γ if there exists a one-to-one correspondence between Γ and Γ^* such that any pseudoline of Γ is a core pseudoline of its corresponding double pseudoline in Γ^*. The two following lemmas are simple consequences of the definitions.

Lemma 6. *The map that assigns to a simple arrangement of pseudolines its set of doubles induces a one-to-one and onto correspondence between the set of isomorphism classes of simple arrangements of pseudolines and the set of isomorphism classes of simple thin arrangements of double pseudolines.* ☐

Lemma 7. *Let* Γ *and* Γ' *be two simple arrangements of pseudolines and let* Γ^* *and* Γ'^* *be double versions of* Γ *and* Γ'. *Assume that* Γ *and* Γ' *are connected by a sequence of two mutations (a merging mutation followed by its 'symmetric' splitting mutation) during which the moving pseudoline is* Γ_i. *Then* Γ^* *and* Γ'^* *are homotopic via a sequence of sixteen mutations during which the only moving double pseudoline is* Γ^*_i. ☐

2.2. **The pumping lemma.** We come now to the statement of our pumping lemma and to the proof of our homotopy theorem.

Lemma 8 (Pumping Lemma)**.** *Let* Γ *be a simple arrangement of double pseudolines, and* $\gamma \in \Gamma$. *Assume that there is a vertex of the arrangement* Γ *lying in the crosscap side of* γ. *Then there is a triangular two-cell of the arrangement* Γ *contained in the crosscap side of* γ *with a side supported by* γ. ☐

Proof. Let $p : \widetilde{\mathcal{P}} \to \mathcal{P}$ be a two-covering of the underlying cross surface $\mathcal{P}$ of Γ and let $\widetilde{\Gamma}$ be the lift of Γ under the covering map p. The lifts of $\gamma \in \Gamma$ under p are denoted γ_+ and γ_-. We note that two curves of $\widetilde{\Gamma}$ have exactly 0 or 2 intersection points depending on whether they are the lifts of the same curve in Γ, or not; by convention if B is one of the two intersection points of two crossing curves of $\widetilde{\Gamma}$ then the other one is denoted B_* (see the top left part of Figure 7 —in this figure the cross surface $\mathcal{P}$ and its two-covering $\widetilde{\mathcal{P}}$ are presented by the polygonal schemes $\alpha\alpha = 1$ and $\alpha\alpha^{-1} = 1$, respectively [35, 32]). Define a γ-*curve* as a maximal subcurve of some $\widetilde{\gamma}' \in \widetilde{\Gamma}$, $\gamma' \neq \gamma$, contained in the cylinder C of $\widetilde{\mathcal{P}}$ bounded by γ_+ and γ_-. We observe that there are two γ-curves per $\widetilde{\gamma}' \in \widetilde{\Gamma}$, $\gamma' \neq \gamma$, and a γ-curve has an endpoint on γ_+ and the other one on γ_-; we denote by $\mathrm{curve}_\gamma(B)$ the γ-curve with endpoint B on γ_+. Two γ-curves $\mathrm{curve}_\gamma(B)$ and $\mathrm{curve}_\gamma(B')$ induce a cell decomposition $C(B, B')$ of the cylinder C which depends only on their number (0, 1 or 2) of intersection points; this cell decomposition is composed of two quadrilaterals or is composed of two triangles and a hexagon or is composed of a digon, two triangles, and an octagon—according to the number 0, 1 or 2 of intersection points—put together as depicted in the bottom part of Figure 7. Define a γ-*triangle* as a triangular face of the cell decomposition of the cylinder C defined by two crossing γ-curves with a side supported by γ_+; the vertex of a γ-triangle not on γ_+ is called its *apex*, the side of a γ-triangle supported by γ_+ called its *base side*, and a γ-triangle is said to be *admissible* if one of its two sides with endpoint its apex is an edge of $\widetilde{\Gamma}$. The interior of the base side of a γ-triangle T is denoted $\mathrm{bs}_\gamma(T)$ and its complement in γ_+ is denoted $\overline{\mathrm{bs}}_\gamma(T)$. Note that $\mathrm{bs}_\gamma(T)$ and $\overline{\mathrm{bs}}_\gamma(T)$ are by definition open intervals. Since by assumption there is a vertex of Γ in the crosscap side of the double pseudoline γ there is a γ-triangle, say $T = ABB'$ with apex A. Let A' be the vertex of $\widetilde{\Gamma}$ that follows B' on the side $B'A$ of T.



$\mathcal{P} : \alpha\alpha = 1$ $\quad$ $\widetilde{\mathcal{P}} : \alpha\alpha^{-1} = 1$

Figure 7. γ-curves.

Then A' is the apex of an admissible γ-triangle $T' = A'B'B''$ with edge side $A'B'$. This proves that there is at least one admissible γ-triangle.

Let now $T = ABB'$ be an admissible γ-triangle with apex A and edge side AB, let A' be the vertex of $\widetilde{\Gamma}$ that follows B' on the side $B'A$ of T, and let $T' = A'B'B''$ be the admissible γ-triangle with apex A' and with edge side $A'B'$. A simple use of the Jordan curve theorem leads to the following lemmas concerning the relative positions of an admissible γ-triangle T with respect to its associated admissible γ-triangle T', possibly in the presence of a third admissible γ-triangle Δ. Figure 8 illustrates these lemmas.

Lemma 9. *Assume that $T = T'$. Then T is a triangular two-cell of $\widetilde{X}$.* $\square$

Lemma 10. *Assume that $T \neq T'$ and that $B'' \in \overline{\mathrm{bs}}_\gamma(T)$. Then*

(1) $\mathrm{curve}_\gamma(B')$ and $\mathrm{curve}_\gamma(B'')$ cross twice (at A' and A'_*) on the side $B'A$ of T,
(2) $\mathrm{bs}_\gamma(T)$ and $\mathrm{bs}_\gamma(T')$ are disjoint,
(3) B'_* and $B''_* \in \overline{\mathrm{bs}}_\gamma(T) \cap \overline{\mathrm{bs}}_\gamma(T')$, and
(4) walking along $\overline{\mathrm{bs}}_\gamma(T) \cap \overline{\mathrm{bs}}_\gamma(T')$ from B'' to B we encounter successively the points B''_* and B'_*. $\square$

Lemma 11. *Assume that $T \neq T'$ and that $B'' \in \mathrm{bs}_\gamma(T)$. Then $\mathrm{curve}_\gamma(B'')$ crosses the side $B'A$ of T exactly once (at A') and $\mathrm{bs}_\gamma(T')$ is contained in $\mathrm{bs}_\gamma(T)$.* $\square$



$B'' \in \overline{\mathrm{bs}}_\gamma(T)$ $B'' \in \mathrm{bs}_\gamma(T)$

$B'' \in \overline{\mathrm{bs}}_\gamma(T)$

FIGURE 8. Relative positions of an admissible γ-triangle T and its derived one T'.

Lemma 12. *Assume that $T \neq T'$, that $B'' \in \overline{\mathrm{bs}}_\gamma(T)$, and that $T = ABB'$ is contained in an admissible γ-triangle $\Delta = XYY'$ with apex X and edge side XY with the property that walking along the base side of Δ from Y to Y' we encounter B' before B (we allow Y' to coincide with B). Then T' is contained in Δ and walking along the base side of Δ from Y to Y' we encounter the point B'' before the point B'.* □

Consider now the sequence of admissible γ-triangles $T_0, T_1, T_2, \dots$ defined inductively by $T_0 = T$ and $T_{k+1} = T'_k$ for $k \geq 0$. A simple combination of Lemmas 10, 11, and 12 leads to the conclusion that the sequence T_k is stationary. According to Lemma 9 the Pumping Lemma follows. □

We are now ready for the proof of our homotopy theorem.

Theorem 2. *Any two arrangements of double pseudolines of the same size and living in the same cross surface are homotopic via a finite sequence of mutations followed by an isotopy; in other terms, mutation graphs are connected.*

Proof. Clearly any arrangement of double pseudolines is homotopic, via a finite sequence of splitting mutations, to a simple one. Now by a repeated application of the Pumping Lemma we see easily that any simple arrangement of double pseudolines is homotopic, via a finite sequence of mutations, to a simple thin one. It remains to use Lemma 6, Lemma 7 and the homotopy theorem for arrangements of pseudolines of Ringel to conclude the proof. ☐

For completeness we mention that one of the standard ways to prove the homotopy theorem for arrangements of pseudolines of Ringel is to show that any arrangement of pseudolines is homotopic, via a finite sequence of mutations followed by an isotopy, to a cyclic arrangement of pseudolines using, in the straightforward manner, *avant la lettre* the following specialization (think of a pair of pseudolines as a pinched double pseudoline) to arrangements of pseudolines of our Pumping Lemma for arrangements of double pseudolines.

Lemma 13 (Pumping Lemma for Arrangements of Pseudolines)**.** *Let Γ be a simple arrangement of pseudolines, let $\gamma, \gamma' \in \Gamma$, $\gamma \neq \gamma'$, and let $M(\gamma, \gamma')$ be one of the two two-cells of the subarrangement $\{\gamma, \gamma'\}$. Assume that there exists a vertex of the arrangement Γ lying in $M(\gamma, \gamma')$. Then there exists a triangular two-cell of the arrangement Γ contained in $M(\gamma, \gamma')$ with a side supported by γ' and a vertex contained in $M(\gamma, \gamma')$.* ☐

Proof. The proof is standard and will not be repeated here; see e.g. [10]. ☐

Remark 1. At this point it is natural to ask if the space of one extensions of an arrangement of double pseudolines is connected under mutations, as is the space of one extensions of an arrangement of pseudolines [25, 2, 44]. A positive answer to that question, providing the key to a practical enumeration algorithm for simple arrangements of at most 5 double pseudolines, is given in [19]. The proof presented in [19] of this connectedness result is based on an enhanced version of the Pumping Lemma which says that, given a double pseudoline γ of an arrangement Γ with the property that the vertices of the arrangement Γ lying on the curve γ are simple, either there are (at least) two fans contained in the crosscap side of the double pseudoline γ with base sides supported by γ or there are no vertices of the arrangement contained in the crosscap side of γ, i.e., γ is thin. The enhanced version of the Pumping Lemma can be easily proved using our theorem saying that any arrangement of double pseudolines is the dual arrangement of a configuration of convex bodies.

2.3. **Enumeration of arrangements.** Representatives of the isomorphism classes of simple arrangements of three double pseudolines are depicted in Figure 9. In this figure each isomorphism class is labeled at its left bottom corner with a symbol to name the arrangement and at its right bottom corner with the order of its automorphism group. The symbol to name an arrangement is composed of the letter C indexed by the 2-sequence of its numbers of 2-cells of size 2 and 3. Such a sequence identifies a unique isomorphism class of arrangements, with one exception: the sequence 25 identifies two isomorphism classes (which have also the same numbers of two-cells of size 4,5,6, etc). To distinguish them we will use the sequences 25_1 and 25_2, where the subscript stands


C_{04} 24
C_{07} 6
C_{18} 4
C_{37} 6
C_{15} 2
C_{43} 2
C_{22} 4
C_{33} 2
C_{32} 2
C_{25} 2
C_{25} 1
C_{36} 12
C_{64} 24

FIGURE 9. Representatives of the thirteen isomorphism classes of simple arrangements of three double pseudolines.

for the order of the automorphism group of the corresponding arrangement. This list was first established by hand, using the connectedness of the corresponding mutation graph. For the numbers of isomorphism classes of simple arrangements on four and five double pseudolines we refer to [19].

Using the exhaustive list of simple arrangements of three double pseudolines we now compute the *martagons* on three and four double pseudolines: martagons are arrangements that play a special rôle in the proof that the isomorphism class of an indexed arrangement of oriented double pseudolines depends only on its chirotope.

An arrangement of $n \geq 3$ double pseudolines Γ is called a *martagon with respect to* a double pseudoline γ of Γ if the vertices of the arrangement on the curve γ are simple and if for any $\gamma' \in \Gamma$, $\gamma' \neq \gamma$, no pair of distinct elements v, v' of

$$\bigcup_{\gamma'' \in \Gamma : \gamma'' \neq \gamma', \gamma} \gamma'' \cap \gamma \tag{2}$$

is separated on the curve γ by a pair of distincts elements u, u' of $\gamma' \cap \gamma$; in other terms: the four intersection points of γ' and γ appear consecutively on the curve γ. For example the arrangements C_{22} and C_{32}, depicted in Figure 9, are martagons with respect to the curved double pseudoline. Figure 10 depicts examples of martagons (with respect to the double pseudoline that do not intersect the dashed pseudoline, red in pdf color) on three and four double pseudolines : in this figure the double pseudolines whose crosscap sides are free of vertices are simply represented by one of their core pseudolines. Observe that the subarrangements of size three of M_1 are C_{22} (3 times) and C_{04}, and those of M_2 are C_{22} and C_{32}, both 2 times. The reader will have no difficulties to add to these examples martagons of arbitrary large size. We leave the verification of the following lemma to the reader.



C_{22} 4 C_{32} 2 M_1 6 M_2 2

FIGURE 10. The martagons on three and four double pseudolines.

Lemma 14. *The only martagons on three and four double pseudolines are the arrangements depicted in Figure 10.* □

3. Geometric representation theorem

In this section we prove the *Geometric Representation Theorem* for double pseudoline arrangements announced in the introduction: any arrangement of double pseudolines is isomorphic to the dual arrangement of a configuration of convex bodies. The main idea of the proof is to show that the property *to be the dual arrangement of a configuration of convex bodies*, defined on the set of arrangements of double pseudolines, is stable under mutations. The main ingredients of the proof are

1. the connectedness of mutation graphs;
2. the coding of the isomorphism class of an arrangement of oriented double pseudolines by its family of *node cycles*;
3. a factorization of the duality transform under the equivalence relation "to have the same isomorphism class of arrangement of bitangents"; such an equivalence class will be termed a *raiponce*;
4. the existence of a projective plane extension for any arrangement of pseudolines [24].

3.1. **Nodes and cycles of an arrangement.** Let Γ be an arrangement of oriented double pseudolines, let I be its indexing set, and let $v(\Gamma)$ be the family of vertices of Γ defined by the three following conditions:

1. the indexing set of $v(\Gamma)$ is the set of unordered pairs ij of (positively or negatively) signed elements of I with $i \neq \pm j$ (we will use both notations $\bar{i}$ and $-i$ for the negative of an index);
2. the $v_\alpha(\Gamma)$, $\alpha \in \{ij, \bar{i}j, i\bar{j}, \overline{ij}\}$, are the four intersection points of the double pseudolines Γ_i and Γ_j;
3. walking along the double pseudoline Γ_i we encounter the $v_\alpha(\Gamma)$, $\alpha \in \{ij, \bar{i}j, i\bar{j}, \overline{ij}\}$, in cyclic order $v_{ij}(\Gamma), v_{\bar{i}j}(\Gamma), v_{i\bar{j}}(\Gamma), v_{\overline{ij}}(\Gamma)$, as illustrated in the left part of Figure 11.



Figure 11. Coding an arrangement of oriented double pseudolines by its cycles.

The reader will easily check that the family $v(\Gamma)$ is well-defined.

The set of *nodes* of Γ, denoted $\mathcal{V}(\Gamma)$, is the quotient of the indexing set of $v(\Gamma)$ under the relation "to index the same vertex of Γ" and the family of *node cycles* of Γ, denoted $\mathcal{C}(\Gamma)$, is the indexed by I family of circular sequences of nodes of Γ that correspond to the circular sequences of vertices of Γ encountered when walking along the oriented double pseudolines of Γ. For example for the non-simple arrangement Υ of three oriented double pseudolines with indexing set $\{1,2,3\}$ depicted in the right part of Figure 11 one has $\mathcal{V}(\Upsilon) = \{A,B,C,D\}$, $\mathcal{C}_1(\Upsilon) = ABCD$, $\mathcal{C}_2(\Upsilon) = ACBD$, and $\mathcal{C}_3(\Upsilon) = ABDC$ where

$$\begin{array}{rcl} A & = & \{12, \overline{13}, \bar{2}3\} \\ B & = & \{\bar{1}2, 13, \overline{23}\} \\ C & = & \{1\bar{2}, \bar{1}3, 2\bar{3}\} \\ D & = & \{\overline{12}, 1\bar{3}, 23\}. \end{array}$$

The family $\mathcal{C}(\Gamma)$ turns out to be a coding of the isomorphism class of Γ.

Theorem 15. *Two arrangements of oriented double pseudolines are isomorphic if and only if they have the same family of node cycles.* $\square$

Proof. Let Γ be an arrangement of oriented double pseudolines, let I be its indexing set, let $\mathcal{F}(\Gamma)$ be the set of flags of the cell poset $X(\Gamma)$ of Γ and let $\sigma_i(\Gamma) : \mathcal{F}(\Gamma) \to \mathcal{F}(\Gamma)$, $i \in \{0,1,2\}$, be its flag operators. The *node, supporting index, orientation* and *side* of a flag $F \in \mathcal{F}(\Gamma)$ are denoted $n(F), i(F), o(F)$, and $s(F)$, and are defined, respectively, as

1. the node of Γ corresponding to the zero-cell of F;
2. the index of the supporting curve of the one-cell of F;
3. the plus sign '+' or the minus sign '-' depending on whether the one-cell of F, endowed with the orientation of its supporting double pseudoline, is outcoming or incoming at the zero-cell of F;
4. the plus sign '+' or the minus sign '-' depending on whether the two-cell of F is contained or not contained in the crosscap side of the supporting double pseudoline of the one-cell of F.

Let $\widehat{\mathcal{F}}(\Gamma) = \{(A, \epsilon, i, \epsilon') \mid i \in I, A \in \mathcal{C}_i(\Gamma), \epsilon, \epsilon' \in \{+,-\}\}$, let $\omega(\Gamma) : \mathcal{F}(\Gamma) \to \widehat{\mathcal{F}}(\Gamma)$ be the (one-to-one and onto) map that assigns to the flag F the 4-uplet $(n(F), o(F), i(F), s(F))$, and, for $i \in \{0,1,2\}$, let

$$\widehat{\sigma}_i(\Gamma) = \omega(\Gamma)\sigma_i(\Gamma)\omega(\Gamma)^{-1}.$$

The 4-uplet $\omega(\Gamma)(F)$ is called the flagcode of F. Figure 12 depicts the first barycentric subdivision of an arrangement of two double pseudolines where each flag is labeled with its flagcode with the convention that $ij{k}^{\epsilon}_{\epsilon'}$ stands for $(ijk, \epsilon, i, \epsilon')$. The table of the operator $\widehat{\sigma}_1(\Gamma)$ in the case where Γ is an arrangement of two double pseudolines is given below

F	ij^-_-	ij^+_-	$\bar{i}j^-_-$	$\bar{i}j^+_-$	$i\bar{j}^-_-$	$i\bar{j}^+_-$	$\overline{ij}^-_-$	$\overline{ij}^+_-$
G	ij^-_+	ij^+_+	$\bar{i}j^-_+$	$\bar{i}j^+_+$	$i\bar{j}^-_+$	$i\bar{j}^+_+$	$\overline{ij}^-_+$	$\overline{ij}^+_+$
$\sigma_1(F)$	ji^-_-	ji^-_+	$j\bar{i}^-_+$	$j\bar{i}^-_-$	$\bar{j}i^+_-$	$\bar{j}i^+_+$	$\overline{ji}^+_+$	$\overline{ji}^+_-$
$\sigma_1(G)$	ji^+_-	ji^+_+	$j\bar{i}^+_+$	$j\bar{i}^+_-$	$\bar{j}i^-_-$	$\bar{j}i^-_+$	$\overline{ji}^-_+$	$\overline{ji}^-_-$

TABLE 1.

Clearly two arrangements of oriented double pseudolines Γ and Γ' are isomorphic if and only if for any $i \in \{0,1,2\}$ the operators $\widehat{\sigma}_i(\Gamma)$ and $\widehat{\sigma}_i(\Gamma')$ coincide. Therefore proving



Figure 12. Flagcodes

our theorem comes down to proving that the operators $\widehat{\sigma}_i(\Gamma)$, $i \in \{0,1,2\}$, depend only on the indexed family $\mathcal{C}(\Gamma)$. Let $\iota : \{+,-\} \to \{+,-\}$ be the transposition that exchanges the plus and minus signs. Clearly

1. (1) $\widehat{\sigma}_2(\Gamma)(A,\epsilon,i,\epsilon') = (A,\epsilon,i,\iota(\epsilon'))$;
2. (2) $\widehat{\sigma}_0(\Gamma)(A,\epsilon,i,\epsilon') = (A',\iota(\epsilon),i,\epsilon')$ where A' is the successor or the predecessor of A in the cycle $\mathcal{C}_i(\Gamma)$ depending on whether ϵ is the plus or minus sign.

Thus it remains to explain why $\widehat{\sigma}_1(\Gamma)$ depends only on $\mathcal{C}(\Gamma)$. (Actually it depends only on $\mathcal{V}(\Gamma)$.) For $J \subseteq I$ with at least two element let $\Gamma|J$ be the restriction of Γ to J and let $i_J : \widehat{\mathcal{F}}(\Gamma|J) \to \widehat{\mathcal{F}}(\Gamma)$ be the induced canonical injection (note that i_J is the identity map on the three last coordinates). For $F \in \widehat{\mathcal{F}}(\Gamma)$, let $U(F)$ be the set of $F_J = i_J\,\widehat{\sigma}_1(\Gamma|J)(i_J)^{-1}(F)$ where J ranges over the set of 2-subsets of I composed of the supporting index i of F and of one of its co-indices, and endow $U(F)$ with the dominance relation $\prec_F$ defined by $F_J \prec_F F_K$ if $F_K = (\widehat{\sigma}_2(\Gamma)(F_J))_{J\Delta K}$ where as usual $J\Delta K$ denotes the set symmetric difference operator. Clearly $\prec_F$ is a total order and $\widehat{\sigma}_1(\Gamma)(F) = \min_{\prec_F} U(F)$. It follows that one can restrict our attention to the case where the size of the set of indices is two. The theorem follows. □

3.2. **Factorization of the duality transform, raiponces.** Recall that a *cyclic* arrangement of pseudolines is a simple arrangement of pseudolines with the property that the maximal size of its two-cells is its number of pseudolines. Figure 13 shows cyclic arrangements of three, four, five and six pseudolines with their two-cells of maximal size marked with little black bullets (red in pdf color). The isomorphism class of a cyclic arrangement of pseudolines depends only of its number of pseudolines; in particular the number of two-cells of maximal size is 4, 3 or 1 depending on whether the size of the arrangement is 3, 4, or larger than 4. Such a two-cell will be called a central cell of the arrangement. We isolate a simple lemma that will be repeatedly used in the sequel.



FIGURE 13. Cyclic arrangements of 3, 4, 5 and 6 pseudolines.

Lemma 16. *Let L be a cyclic arrangement of pseudolines, let ∇ be a central cell of L and let A be a vertex of ∇ that is an intersection of the pseudolines x and y of L. Let z be a pseudoline such that (1) z is tangent to ∇ at A and (2) the family $L' = L \setminus \{x, y\} \cup z$ is a simple arrangement of pseudolines. Then (1) L' is cyclic and (2) ∇ is contained in a central two-cell ∇' of L'.* □

Let I be a finite set of indices and let $\mathcal{N}(I)$ be the set of unordered pairs ij of signed elements of I with $i \neq \pm j$. By definition a *raiponce* L on I is a simple arrangement of pseudolines indexed by $\mathcal{N}(I)$ such that

(1) for any $i, j \in I$, $i \neq j$, the subarrangement of L_α, $\alpha \in \{ij, \bar{i}j, i\bar{j}, \bar{i}\bar{j}\}$, is an arrangement of four pseudolines; we denote by $\nabla_{ij}(L)$ its unique oriented two-cell such that walking along its boundary we encounter the pseudolines L_α, $\alpha \in \{ij, \bar{i}j, i\bar{j}, \bar{i}\bar{j}\}$, in the circular order $L_{ij}, L_{\bar{i}j}, L_{i\bar{j}}, L_{\bar{i}\bar{j}}$, as illustrated in the left part of Figure 14; note that $\nabla_{ij}(L)$ and $\nabla_{ji}(L)$ are by construction disjoint and that their closures share a vertex but no edge;

(2) for any $i \in I$ the subarrangement of L_α, $j \in I \setminus i$, $\alpha \in \{ij, \bar{i}j, i\bar{j}, \bar{i}\bar{j}\}$, is cyclic and walking along the boundary of one of its oriented central cells we encounter for any $j \in I \setminus i$ the pseudolines L_α, $\alpha \in \{ij, \bar{i}j, i\bar{j}, \bar{i}\bar{j}\}$, in the circular order $L_{ij}, L_{\bar{i}j}, L_{i\bar{j}}, L_{\bar{i}\bar{j}}$; this oriented central cell is necessarily the intersection of the $\nabla_{ij}(L)$, $j \in I \setminus i$, and will be denoted $\nabla_i(L)$.

Let $\mathcal{V}(L)$ be the quotient of $\mathcal{N}(I)$ under the relation "to index the same pseudoline of L" and let $\mathcal{C}(L)$ be the family indexed by I of circular sequences of elements of $\mathcal{V}(L)$ encountered when walking along the (oriented) boundaries of the polygons $\nabla_i(L)$. An element of $\mathcal{C}(L)$ will be called a *cycle* of L. The reader will easily check that the family of cycles of the raiponce L^0 depicted in the right part of Figure 14 coincides with the



FIGURE 14.

family of cycles of the arrangement of oriented double pseudolines Υ depicted in the right part of Figure 11. Let now Δ be a finite family of pairwise disjoint oriented convex bodies of a projective plane whose family of bitangents is simple and let Δ^* be its dual (oriented) arrangement. Clearly the family $v(\Delta^*)$ of vertices of Δ^*—see Section 3.1 for its definition—is a raiponce on I. The following lemma claims that any raiponce is the arrangement of bitangents of a family of convex bodies and that the map $\Delta \mapsto v(\Delta^*)$ is compatible with the duality transform. The proof is easy.

Lemma 17. *Let L be a raiponce on I, let $\mathcal{G}$ be a projective plane extension of the underlying arrangement of pseudolines of L, and let $\mathcal{R}(L,\mathcal{G})$ be the class of finite families Δ of pairwise disjoint oriented convex bodies of $\mathcal{G}$ indexed by I such that for any $i \in I$*

1. *Δ_i is inscribed in the polygon $\nabla_i(L)$, and*
2. *the orientations of Δ_i and $\nabla_i(L)$ are coherent.*

Then

1. *$\mathcal{R}(L,\mathcal{G})$ is nonempty, and*
2. *for any $\Delta \in \mathcal{R}(L,\mathcal{G})$, the isomorphism class of Δ^* depends only on L.* □

Proof. The first point is clear since by construction the closures of the $\nabla_i(L)$ intersect pairwise in at most one vertex. Similarly the second point is clear since by construction $\mathcal{V}(\Delta^*) = \mathcal{V}(L)$ and $\mathcal{C}(\Delta^*) = \mathcal{C}(L)$. □

An element of $\bigcup_{\mathcal{G}} \mathcal{R}(L,\mathcal{G})$, where $\mathcal{G}$ ranges over the set of projective plane extensions of the underlying arrangement of L, will be termed a completion of L, and a raiponce L will be said to be a primal representation of an arrangement of double pseudolines Γ if the isomorphism class of Γ is the isomorphism class of the completions of L. For example the raiponce L^0 depicted in the right part of Figure 14 is a primal representation of the arrangement of three double pseudolines Υ depicted in the right part of Figure 11. According to the previous discussion the properties 'to be the dual arrangement of a family of pairwise disjoint convex bodies' and 'to have a primal representation' are equivalent. The next step is devoted to the proof that this last property is stable by mutation.

3.3. **Stability by mutations.**

Theorem 18. *The property 'to have a primal representation' is stable by mutation.* ☐

Before embarking on the proof we isolate two simple properties of primal representations. The proofs are easy.

Lemma 19. *Let L be a primal representation of an arrangement of double pseudolines Γ indexed by I, let x be a vertex of Γ, and let $[x]$ be its corresponding pseudoline in L. Then for any index $i \in I$*

(1) x belongs to the crosscap side of Γ_i if and only if $[x]$ pierces $\nabla_i(L)$;
(2) x belongs to Γ_i if and only if $[x]$ is a supporting pseudoline of $\nabla_i(L)$;
(3) x belongs to the disk side of Γ_i if and only if $[x]$ avoids the closure of $\nabla_i(L)$.

as illustrated in Figure 15. ☐

Proof. This is a direct application of Theorem 1 applied to any completion Δ of L such that for index i one has $\Delta_i = \nabla_i(L)$. ☐



FIGURE 15. Illustration of Lemma 19

Lemma 20. *Let L be a primal representation of an arrangement of double pseudolines Γ indexed by I, let σ be a one-cell of Γ, and let $[x]$ and $[y]$ be the oriented pseudolines of L corresponding to the endpoints of any lift $\widetilde{\sigma}$ of σ in a two-covering of the underlying cross surface of Γ. Then*

(1) σ is contained in the crosscap side of Γ_i if and only if $\nabla_i(L)$ is transversal to the non-cyclic two-cell of the 2-arrangement $[x], [y]$ as depicted in the three circular diagrams of the second row of the top part of Figure 16;
(2) σ is contained in the topological closure of the disk side of Γ_i if and only if $\nabla_i(L)$ is contained in the topological closure of the cyclic two-cell of the 2-arrangement $[x], [y]$ as depicted in the four circular diagrams of the bottom part of Figure 16. ☐



FIGURE 16. Illustration of Lemma 20

Proof of Theorem 18. Let L be a primal representation of an arrangement of oriented double pseudolines Γ, and consider a mutation connecting Γ to an arrangement Γ'. Our goal is to show that Γ' has a primal representation L'. Without loss of generality one can assume that Γ is the dual arrangement of a completion Δ of L.

We first examine the case of a merging mutation.

Let Σ be the complex of adjacent triangular two-cells of Γ involved in the merging mutation and let $\widetilde{\Sigma}$ be one of its two lifts in a two-covering of the underlying cross surface. We consider the set of vertices of $\widetilde{\Sigma}$ as an arrangement Ψ of oriented pseudolines and we introduce the one-level ℓ of the subarrangement Ψ_0 composed of the three vertices of the boundary $\partial\widetilde{\Sigma}$ of $\widetilde{\Sigma}$ with respect to its unique two-cell σ_0 with cyclic boundary; note that ℓ is by construction a pseudoline and that any pseudoline in L not in Ψ cross ℓ in at most three points. Figure 17 depicts the complex Σ of a merging mutation, the subarrangement Ψ_0 with its cyclic two-cell σ_0 marked, and the one-level ℓ of Ψ_0 with respect to σ_0.



FIGURE 17. The complex Σ of triangular two-cells involved in the merging mutation connecting Γ to Γ', the arrangement Ψ_0 composed of the vertices of the boundary of the complex $\widetilde{\Sigma}$, and the one-level ℓ of the arrangement Ψ_0.

Let I be the indexing set of Γ and let L' be the simple family of pseudolines indexed by $\mathcal{N}(I)$ defined by

$$L'_\tau = \begin{cases} \ell & \text{if } L_\tau \text{ is a vertex of } \Sigma; \\ L_\tau & \text{otherwise.} \end{cases} \tag{3}$$

Lemma 21. *We claim that*

1. *L' is a simple arrangement of pseudolines;*
2. *L' is a primal representation of the arrangement Γ'.*

Proof. Let K be the set of indices of the supporting double pseudolines of the one-cells of Σ, let $K' \subseteq K$ be the set of three indices of the three supporting double pseudolines of the three sides of the boundary of Σ, and let $w \in K'$ be the index of the moving curve of the mutation. We denote by $\widehat{w}$ the vertex of the boundary of Σ opposite to the side supported by Γ_w, and for any $t \in K \setminus \{w\}$ we denote by $\widehat{t}$ the vertex of Σ intersection of the double pseudolines Γ_t and Γ_w. Let Ψ_0^+ be the arrangement Ψ_0 augmented with the line $\widehat{t}$ if $t \in K \setminus K'$; the arrangement Ψ if $t = w$; the arrangement Ψ_0 otherwise. We denote by L^* the sub-raiponce of L obtained by deleting the L_{ijk} with $i \in K \setminus K'$. Finally let $\widehat{u}_1, \widehat{u}_2, \ldots, \widehat{u}_m$ be the sequence of vertices $\neq \widehat{w}$ of Σ ordered along Γ_w.

Applying Lemma 20 to the one-cells of $\widetilde{\Sigma}$ (and using induction on the size of $K \setminus K'$) we see easily that the relative position of $\nabla_t(L)$ in the arrangement Ψ_0^+ depends only on whether the triangular two-cells of Σ are contained in the disk side $\mathcal{D}(\Gamma_t)$ of Γ_t or contained in crosscap side $\mathcal{M}(\Gamma_t)$ of Γ_t, as indicated in Figure 18. Furthermore one can also check that

1. for any $t \in K \setminus \{w\}$ the pseudoline ℓ is tangent to $\nabla_t(L)$ at the intersection point of $\widehat{w}$ and $\widehat{t}$;
2. the pseudolines in $\Psi \setminus \Psi_0$ cross the pseudoline ℓ all in three points or all in one point;
3. the arrangement Ψ is cyclic;
4. the relative position of $\nabla_w(L^*)$ in the arrangement Ψ_0 depends only on whether the triangular two-cells of Σ are contained in $\mathcal{D}(\Gamma_w)$ or in $\mathcal{M}(\Gamma_w)$ as indicated in Figure 18; in particular we note that ℓ is tangent to $\nabla_w(L^*)$ at the intersection point of $\widehat{u}_1$ and $\widehat{u}_m$.

Pick now a pseudoline ℓ' such that $L \cup \{\ell'\}$ is a simple arrangement of pseudolines. Assume that ℓ' and ℓ cross three times. Clearly ℓ' avoids the cyclic two-cell of Ψ_0 and



FIGURE 18. Relative position of $\nabla_t(L)$ in the arrangement Ψ_0.

consequently—thanks to our previous discussion on the position of the $\nabla_t(L)$ in the arrangement Ψ_0^+—ℓ' is transversal to any $\nabla_t(L)$, $t \in I \setminus (K \setminus K')$, not contained in σ_0. It follows that $\ell' \notin L \setminus \Psi$ and, consequently, there is no pseudoline of $L \setminus \Psi$ crossing ℓ three times: thus L' is a simple arrangement of pseudolines.

We now prove that L' is a raiponce and a primal representation of Γ'. Given a subfamily S of L we set S' to be the corresponding subfamily of L'. For any index $i \in I$ let M_i be the arrangement of pseudolines L_α, $\alpha \in \{ij, \bar{i}j, i\bar{j}, \overline{ij}\}$, $j \in I \setminus \{i\}$, and let $N_{ij} = M_i \cap M_j$, $i \neq j \in I$. Observe that N_{ij} contains at most one element of Σ and that $\ell \notin L$: consequently N'_{ij} is an arrangement of four pseudolines. By construction

$$M'_t = \begin{cases} M_t \setminus \{\widehat{w}, \widehat{t}\} \cup \{\ell\} & \text{if } t \in K \setminus \{w\}; \\ M_t \setminus \Sigma \cup \{\ell\} & \text{if } t = w; \\ M_t & \text{otherwise.} \end{cases} \tag{4}$$

Since for any $i \in K \setminus \{w\}$ the pseudoline ℓ is tangent to $\nabla_i(L)$ at the intersection point of $\widehat{w}$ and $\widehat{t}$, and since ℓ is tangent to $\nabla_w(L^*)$ at the intersection point of $\widehat{u}_1$ and $\widehat{u}_m$

it follows, according to Lemma 16, that for any i the arrangement M'_i is cyclic, that $\nabla_i(L)$ is contained in one of its central two-cell ∇'_i, and that walking along its boundary (oriented according to the orientation of $\nabla_i(L)$) we encounter for any $j \in I \setminus i$ the pseudolines L'_α, $\alpha \in \{ij, \bar{i}j, i\bar{j}, \overline{ij}\}$, in the circular order $L'_{ij}, L'_{\bar{i}j}, L'_{i\bar{j}}, L'_{\overline{ij}}$; consequently L' is a raiponce and is (by construction) a primal representation of Γ'. $\square$



$\widehat{w} \in \mathcal{D}(\Gamma'_w)$ $\widehat{w} \in \mathcal{M}(\Gamma'_w)$

FIGURE 19. The complex Σ of triangular two-cells involved in the merging mutation connecting Γ to Γ', the arrangement Ψ_0 composed of the vertices of the boundary of the complex $\widetilde{\Sigma}$, and the one-level ℓ of the arrangement Ψ_0.

We now examine the case of a splitting mutation.

Let K be the set of indices of the double pseudolines involved in the splitting mutation, let w be the index of the moving double pseudoline, let $\widehat{w}$ be the vertex involved in the mutation, and for any $v \in K \setminus \{w\}$ let $x(v) \in \{wv, \overline{w}v, w\overline{v}, \overline{wv}\}$ defined by the condition $L_{x(v)} = \widehat{w}$.

Let w^* be a double pseudoline containing $\widehat{w}$ in its Möbius strip such that any pseudoline of L crosses w^* in exactly two points and such that no vertex of the arrangement L belongs to the Möbius strip $\mathcal{M}(w^*)$ bounded by w^*. The pseudolines of L induce a decomposition of $\mathcal{M}(w^*)$ into quadrilateral regions. In particular the trace of the central cell of the raiponce L indexed by w onto $\mathcal{M}(w^*)$ is one of its quadrilateral regions that we shall denote by Q. We denote by S and S^* the sides of Q supported by $\widehat{w}$ and w^*, respectively, and we denote by Q' the second quadrilateral region of $\mathcal{M}(w^*)$ bounded by S.

Let B_1 be a generic point of Q if $\widehat{w} \in \mathcal{D}(\Gamma'_w)$; otherwise let B_1 be a generic point of Q'.

For any $i \in K \setminus \{w\}$ we insert a generic point B_i on the interior of the edge of the central cell of L indexed by i supported by $\widehat{w}$, and we insert in the underlying pseudoline arrangement of L a pseudoline ℓ_i such that

(1) ℓ_i goes through the points B_1 and B_i, and is contained in $\mathcal{M}(w^*)$;
(2) the vertices of $L \cup \Psi$ are simple, excepted B_1;

and we perturb the pencil of pseudolines ℓ_i in the vicinity of B_1 into a cyclic arrangement ℓ_i^* with a central cell containing S^* or S depending on whether $\widehat{w} \in \mathcal{D}(\Gamma'_w)$ or not.

Let now L' be the simple family of pseudolines indexed by $\mathcal{N}(I)$ defined by

$$L'_\tau = \begin{cases} \ell_v^* & \text{if } \tau = x(v) \text{ with } v \in K \setminus \{w\}; \\ L_\tau & \text{otherwise.} \end{cases}$$

A simple case analysis shows that L' is a well-defined raiponce and is a primal representation of Γ'. Details are left to the reader. $\square$

4. Cycles, cocycles and chirotopes

In this section we prove that (1) the isomorphism class of an indexed arrangement of oriented double pseudolines depends only on its family of isomorphism classes of subarrangements of size three, i.e., depends only on what we have called its chirotope; and that (2) the duality map is compatible with the isomorphism relations on the set of configurations of convex bodies and on the set of arrangements of double pseudolines and induces a one-to-one correspondence between the set of isomorphism classes of indexed configurations of oriented convex bodies and the set of isomorphism classes of indexed arrangements of oriented double pseudolines. The main ingredients of our proof are

1. the coding of the isomorphism class of an indexed arrangement of oriented double pseudolines by its family of *side cycles*, which are useful variants of the node cycles of Section 3;
2. the list of martagons on three and four double pseudolines, established in Section 2;
3. the injectivity of cocycle maps of families of two pairwise disjoint convex bodies.

4.1. **Side cycles.** Side cycles are more homogeneous variants of the node cycles of Section 3 that do not encode explicitly the nodes of the arrangement at the cost of doubling the number of cycles : there are two side cycles per double pseudoline : one cycle of *disk type* and the other one of *crosscap type*. Let Γ be an indexed arrangement of oriented double pseudolines and let I be its indexing set. The *side cycle of disk type* of the double pseudoline Γ_i or indexed by i, denoted D_i, is the circular sequence of indices of the double pseudolines crossed by the side wheel of a sidecar rolling on Γ_i, side wheel on the disk side of Γ_i, each index being signed positively or negatively depending on whether the crossed double pseudoline is (locally) oriented away or towards Γ_i. Similarly the *side cycle of crosscap type* of the double pseudoline Γ_i, denoted M_i, is the circular sequence of indices of the double pseudolines crossed by the side wheel of a sidecar rolling on Γ_i, side wheel on the crosscap side of Γ_i, each index being signed positively or negatively depending on whether the crossed double pseudoline is (locally) oriented away or towards Γ_i. Observe that for simple arrangements the side cycles of disk type and crosscap type are reverse to one another.

Example 3. The side cycles of disk type of an arrangement Γ on two double pseudolines, say indexed by i, j, are

$$\begin{array}{rl} i: & \overline{jj}jj \\ j: & \overline{ii}ii. \end{array}$$

as illustrated in Figure 20 where we have displayed the first barycentric subdivision of the one-skeleton of the arrangement and labeled each edge of the subdivision with the index of its supporting curve signed positively or negatively depending on whether the edge is outcoming or incoming at the vertex of the arrangement to which it is incident. Observe that each symbol in these cycles corresponds in the natural way to a unique node of the arrangement, namely the linear sequence $\overline{jj}jj$ corresponds to the linear sequence of nodes $\{ij\}\{\bar{i}j\}\{i\bar{j}\}\{\overline{ij}\}$, as illustrated in Figure 20. The side cycles of crosscap type of Γ coincide with its side cycles of disk type but now the linear sequence $\overline{jj}jj$ corresponds to the linear sequence of nodes $\{i\bar{j}\}\{\overline{ij}\}\{ij\}\{\bar{i}j\}$.



FIGURE 20. Side cycles of an arrangement of two double pseudolines.

Example 4. The side cycles of disk type and crosscap type of the arrangement depicted in the left part of Figure 21 are

$$\begin{array}{ll} 1: & 32\overline{2}3\overline{32}2\overline{3} \\ 2: & \overline{3}1\overline{13}3\overline{1}13 \\ 3: & \overline{21}1\overline{2}21\overline{1}2 \end{array} \qquad \text{and} \qquad \begin{array}{ll} 1: & \overline{233}2233\overline{2} \\ 2: & \overline{1}3311\overline{331} \\ 3: & 122\overline{1122}1. \end{array}$$

Similarly the side cycles of disk type and crosscap type of the arrangement depicted in the right part of Figure 21 (obtained from the left arrangement by a splitting mutation) are

$$\begin{array}{ll} 1: & 32\overline{2}3\overline{32}2\overline{3} \\ 2: & \overline{3}1\overline{13}3\overline{1}13 \\ 3: & \overline{21}1\overline{2}21\overline{1}2 \end{array} \qquad \text{and} \qquad \begin{array}{ll} 1: & \overline{323}2233\overline{2} \\ 2: & 3\overline{1}311\overline{331} \\ 3: & 212\overline{1122}1. \end{array}$$

Note that these two arrangements have the same side cycles of disk type but differ in their side cycles of crosscap type.



FIGURE 21. Two arrangements with the same side cycles of disk type.

Let Γ be an indexed arrangement of oriented double pseudolines with indexing set I, let D_i be its side cycles of disk type and M_i those of crosscap type. Let S_i be the result of replacing in D_i the linear subsequences $\overline{jj}jj$, $j \neq i$, by the linear sequences $\{ij\}\{\bar{i}j\}\{i\bar{j}\}\{\overline{ij}\}$; similarly let T_i be the result of replacing in M_i the linear subsequences $jj\overline{jj}$, $j \neq i$, by the linear sequences $\{ij\}\{\bar{i}j\}\{i\bar{j}\}\{\overline{ij}\}$. Clearly there is one-to-one correspondence between the vertices of the arrangement lying on the curve indexed by i and the maximal factors $\{i_1j_1\}\{i_2j_2\}\{i_3j_3\}\ldots\{i_kj_k\}$ of S_i with $j_l \neq \pm j_{l'}$ for all $1 \leq l < l' \leq k$ that appear in reverse order $\{i_kj_k\}\ldots\{i_3j_3\}\{i_2j_2\}\{i_1j_1\}$ in T_i, *prime factors* for short. More precisely

(1) the node of vertex v associated with the prime factor $\{i_1j_1\}\{i_2j_2\}\{i_3j_3\}\ldots\{i_kj_k\}$ of S_i is the set of $\{i_lj_l\}\otimes\{i_{l'}j_{l'}\}$, $1 \leq l \leq l' \leq k$, where $i_lj_l \otimes i_{l'}j_{l'}$ is the element of the 4-set $\{j_lj_{l'}, \overline{j_l}j_{l'}, j_l\overline{j_{l'}}, \overline{j_lj_{l'}}\}$ indexing the intersection point of Γ_{j_l} and $\Gamma_{j_{l'}}$ that coincides with v; as illustrated in Figure 22—which depicts implicitly the 4×4 possible nodes involving three curves indexed by i, j and j_* where $i \in I$,$j, j_* \in \pm I$; the dashed sides stand for the crosscap sides—this element depends only on the sole information contains in the (ordered) pair $\{i_lj_l\}\{i_{l'}j_{l'}\}$ and the multiplication table of $\otimes$ is the following

$$\begin{cases} ij \otimes ij & = & ji \\ ij \otimes ij_* & = & j\overline{j_*} \\ ij \otimes \bar{i}j_* & = & \overline{jj_*} \\ \bar{i}j \otimes ij_* & = & jj_* \\ \bar{i}j \otimes \bar{i}j_* & = & \bar{j}j_* \end{cases}$$

where $i \in I$, $j, j_* \in \pm I$ with $j \neq \pm j_*$; and conversely

(2) the prime factor of S_i corresponding to the vertex v is the sequence of $\{\eta j\}$, $\eta = \pm i$, that belongs to the node N of v, ordered according to the dominance relation $\{\eta j\} \prec \{\eta' j'\}$ if $\{\eta j\} \otimes \{\eta' j'\} \in N$.



$\{ij\}\{ij_*\}$ $\{ij\}\{\bar{i}j_*\}$ $\{\bar{i}j\}\{ij_*\}$ $\{\bar{i}j\}\{\bar{i}j_*\}$

Figure 22.

Thus node cycles and side cycles carry exactly the same information about the arrangement. Since, according to Theorem 15, two indexed and oriented arrangements are isomorphic if and only if they have the same family of node cycles we get

Theorem 22. *Two indexed arrangements of oriented double pseudolines are isomorphic if and only if they have the same family of side cycles.* □

Let X be an arrangement of double pseudolines, let X_* be an indexed and oriented version of X, say indexed by the indexing set I, and extend X_* to the set of negatives of the elements of I by assigning to the negative of the index i the reoriented version of the double pseudoline assigned to i. Let G_I be the group of permutations of the signed indices which are compatible with the operation of taking the negative of an index, let G_{X_*} be the stabilizer of X_*, i.e., the subgroup of G_I whose elements are the permutations σ such that $X_*\sigma$ and X_* are isomorphic, and let G_X be the group of automorphisms of X, that we think as a subgroup of the group of permutations of the signed (or oriented) double pseudolines of the arrangement. Clearly the map that assigns to $\sigma \in G_X$ its conjugate $X_*^{-1}\sigma X_* \in G_I$ under X_* is a monomorphism of G_X onto G_{X_*}. Thus we can see G_X as a subgroup G_{X_*} of G_I and the number of distinct indexed and oriented versions of X is the index $[G_I : G_{X^*}]$ of G_{X_*} in G_I. In the sequel we use the notation $X(\sigma)$ for the arrangement $X_*\sigma$, $\sigma \in G_I$; hence $X(1) = X_*$, where 1 is the unit of G_I.

Example 5. Figure 23 depicts an arrangement Z on 3 double pseudolines and two of its indexed and oriented versions on the indexing set $\{1, 2, 3\}$. The group G_Z is, as explained in Section 1, S_4. Thus the number of distinct indexed and oriented versions of Z is $3!2^3/24 = 2$. The group G_{Z_*} is of order 24 generated by the permutations $\overline{1}32$ and $\overline{12}3$ and $2\overline{31}$ (which correspond to the automorphisms τ_1, τ_2 and τ_3 of Section 1), respectively. Its two cosets are

$$G_{Z_*} = \left\{\begin{matrix} 123 & 231 & 312 \\ \overline{12}3 & \overline{23}1 & \overline{31}2 \\ \overline{1}2\overline{3} & \overline{2}3\overline{1} & \overline{3}1\overline{2} \\ 1\overline{23} & 2\overline{31} & 3\overline{12} \\ 21\overline{3} & 32\overline{1} & 13\overline{2} \\ 3\overline{2}1 & 1\overline{3}2 & 2\overline{1}3 \\ \overline{1}32 & \overline{2}13 & \overline{3}21 \\ \overline{213} & \overline{321} & \overline{132} \end{matrix}\right\}, \qquad (213)G_{Z_*} = \left\{\begin{matrix} 213 & 321 & 132 \\ \overline{21}3 & \overline{32}1 & \overline{13}2 \\ \overline{2}1\overline{3} & \overline{3}2\overline{1} & \overline{1}3\overline{2} \\ 2\overline{13} & 3\overline{21} & 1\overline{32} \\ 12\overline{3} & 23\overline{1} & 31\overline{2} \\ 3\overline{1}2 & 1\overline{2}3 & 2\overline{3}1 \\ \overline{2}31 & \overline{3}12 & \overline{1}23 \\ \overline{123} & \overline{231} & \overline{312} \end{matrix}\right\}.$$



Figure 23.

Example 6. As illustrated in Figure 24 the two families of cycles on the indexing set $1, 2, 3, 4$

$$\begin{array}{ll} 1: & 22\overline{22}33\overline{33}44\overline{44} \\ 2: & 441\overline{133441}133 \\ 3: & 221\overline{144221}144 \\ 4: & 331\overline{122331}122 \end{array} \quad \text{and} \quad \begin{array}{ll} 1: & \overline{22}22\overline{33}33\overline{4}44\overline{4} \\ 2: & \overline{33}1\overline{14}433\overline{1}14\overline{4} \\ 3: & \overline{22}4\overline{4}1\overline{1}22\overline{4}4\overline{1}1 \\ 4: & \overline{22}1\overline{1}2233\overline{1}1\overline{33} \end{array}$$

are the side cycles of disk type of indexed and oriented versions of the two martagons M_1 and M_2 on four double pseudolines, depicted in Figure 10. The automorphism group of M_1 is S_3 (dihedral group of order 6) generated by the involutions $\overline{1324}$ and $\overline{1243}$, for example. The automorphism group of M_2 is of order 2 generated by the permutation $\overline{1324}$.



$M_1(1234)$ $4!2^4/6$ $M_2(1234)$ $4!2^4/2$

FIGURE 24. Oriented and indexed versions of the martagons on 4 double pseudolines.

We close this section by introducing, in Figures 25 and 26, one oriented and indexed version of each of the thirteen simple arrangements on three double pseudolines of Figure 9. Each diagram is labeled at its right top corner by its number of (distinct) reindexed and reoriented versions and at its right bottom corner by its side cycles of disk type. We let the reader check that the four subarrangements of size three of $M_1(1234)$ are $C_{22}(1\overline{23})$, $C_{22}(1\overline{34})$, $C_{22}(1\overline{42})$ and $C_{04}(234)$. Similarly the four subarrangements of size three of the martagon $M_2(1234)$ are $C_{22}(123)$, $C_{22}(42\overline{3})$, $C_{32}(142)$, $C_{32}(143)$.


2
1
3
2
$C_{04}(123)$
$2233\overline{2233}$
$3311\overline{3311}$
$1122\overline{1122}$
8
3
1
3
2
$C_{07}(123)$
$2233\overline{2323}$
$3131\overline{3311}$
$112\overline{1}2\overline{122}$
12
2
3
1
$C_{18}(123)$
$\overline{3}233\overline{232}2$
$\overline{1}131\overline{331}3$
$\overline{2}12\overline{1}2\overline{12}1$
8
1
2
3
$C_{37}(123)$
$\overline{3}23\overline{2}3\overline{32}2$
$\overline{1}131\overline{313}3$
$\overline{2}21\overline{1}2\overline{12}1$
2
2
3
1
$C_{15}(123)$
$33\overline{223}22\overline{3}$
$3\overline{1}31\overline{3}1\overline{31}$
$\overline{1}1\overline{22}122\overline{1}$
24
2
3
1
$C_{43}(123)$
$2\overline{3}3\overline{2}3\overline{32}2$
$\overline{1}13\overline{3}1\overline{13}3$
$\overline{2}21\overline{1}2\overline{1}1\overline{2}$

FIGURE 25. Indexed and oriented versions of the arrangements $C_{04}, C_{07}, C_{18}, C_{37}, C_{15}, C_{43}$.



$C_{22}(123)$

$\overline{33}33\overline{22}22$
$33\overline{1}1\overline{33}1\overline{1}$
$22\overline{1}1\overline{22}1\overline{1}$



$C_{33}(123)$

$2\overline{3}3\overline{2}3\overline{2}2\overline{3}$
$33\overline{1}1\overline{3}1\overline{13}$
$\overline{1}2\overline{1}1\overline{22}21$



$C_{32}(123)$

$\overline{3}33\overline{322}22$
$3\overline{1}13\overline{3}1\overline{13}$
$22\overline{1}1\overline{22}1\overline{1}$



$C_{25_2}(123)$

$2\overline{3}33\overline{22}2\overline{3}$
$33\overline{1}1\overline{3}1\overline{31}$
$\overline{1}2\overline{1}1\overline{22}12$



$C_{25_1}(123)$

$32\overline{2}3\overline{2}2\overline{33}$
$1\overline{313}33\overline{1}1$
$\overline{12}1\overline{2}21\overline{1}2$



$C_{36}(123)$

$3\overline{2}3\overline{3}2\overline{3}2\overline{2}$
$3131\overline{1313}$
$1\overline{1212}212$



$C_{64}(123)$

$2\overline{2}3\overline{32}2\overline{3}3$
$3\overline{3}1\overline{13}3\overline{1}1$
$1\overline{1}2\overline{21}1\overline{2}2$

FIGURE 26. Indexed and oriented versions of the arrangements $C_{22}, C_{33}, C_{32}, C_{25_2}, C_{25_1}, C_{36}, C_{64}$.

4.2. **Chirotopes.** We are now ready to show that the isomorphism class of an arrangement of oriented double pseudolines depends only on its chirotope; cf. first part of Theorem 4.

Theorem 23. *The map that assigns to an isomorphism class of indexed arrangements of oriented double pseudolines is chirotope is one-to-one.* □

Proof. Let Γ be an indexed arrangement of oriented double pseudolines. According to Theorem 15 the isomorphism class of Γ depends only on the family of cycles of Γ; therefore it is sufficient to show that the family of cycles of Γ depends only on the chirotope of Γ.

Clearly the set of nodes of Γ depends only on the chirotope of Γ, and clearly one can restrict our attention to the case where Γ has four elements, say $\Gamma_1, \Gamma_2, \Gamma_3$ and Γ_4.

We now show that the cycle of Γ indexed by 1 depends only on the chirotope of Γ. We write $\mathcal{N}(k)$, $k \in \{2,3,4\}$, for the set of nodes of Γ indexing the intersection points of Γ_1 and Γ_k, $\mathcal{N}(k,k',\ldots)$ for $\mathcal{N}(\{k,k',\ldots\})$, and for any $A, A' \in \mathcal{N}(2,3,4)$, $A \neq A'$, we write $[A, A']$ for the set of $X \in \mathcal{N}(2,3,4)$, including A and A', that appear between A and A' on the cycle of Γ indexed by 1.

Let $A, A' \in \mathcal{N}(2)$, $A \neq A'$, and let $B, B' \in \mathcal{N}(2,3,4)$. We say that the pair (A, A') separates the pair (B, B') if $B \in [A, A']$ and $B' \in [A', A]$. Clearly one can decide, using only the chirotope of Γ,

(1) whether B belongs to the interval $[A, A']$ or not; and
(2) whether the pair (A, A') separates the pair (B, B') or not.

Assume that a pair (A, A') of distinct elements of $\mathcal{N}(2)$ separates a pair (B, B') of elements of $\mathcal{N}(3,4)$. (In particular this happens if one of the intersection points between Γ_1 and Γ_2 is a non-simple vertex of the arrangement Γ.) In that case a pair X, Y of distinct elements of $\mathcal{N}(3,4)$ lying in the open interval $[A, A'] \setminus \{A, A'\}$ appears in the linear order XY in the interval $[A, A']$ if and only if B', X, Y appear in the cyclic order $B'XY$ on the cycle of the arrangement $\{\Gamma_1, \Gamma_3, \Gamma_4\}$ indexed by 1. Consequently the cycle of Γ indexed by 1 depends only on the chirotope of Γ and we are done. Similarly we are done if a pair of distinct elements of $\mathcal{N}(3)$ separates a pair of elements of $\mathcal{N}(2,4)$, or if a pair of distinct elements of $\mathcal{N}(4)$ separates a pair of elements of $\mathcal{N}(2,3)$. Thus it remains to examine the case where for every $k \in \{2,3,4\}$ no pair of distinct elements of $\mathcal{N}(k)$ separates a pair of elements of $\mathcal{N}(\{2,3,4\} \setminus \{k\})$, i.e., using the terminology introduced in the previous section, the case where the arrangement Γ is a martagon with respect to Γ_1. According to Lemma 14 and the notations introduced in Example 6, this means that, up to permutation of the indices $1, 2, 3, 4$ and their negatives, $\Gamma = M_1(1234)$ or $M_2(1234)$. The theorem follows. Indeed if the chirotope of Γ is the chirotope of $M_1(1234)$ then the family of side cycles (of disk type) of Γ is necessarily either the family

$$\begin{array}{ll} \mathcal{C}_1: & 22\overline{22}33\overline{33}44\overline{44} \\ \mathcal{C}_2: & 441\overline{133441}133 \\ \mathcal{C}_3: & 221\overline{144221}144 \\ \mathcal{C}_4: & 331\overline{122331}122 \end{array}$$

of side cycles of $M_1(1234)$ or the family

$$\begin{array}{ll} \mathcal{C}_1^* : & 33\overline{33}22\overline{22}44\overline{44} \\ \mathcal{C}_2^* : & 441\overline{133441}133 \\ \mathcal{C}_3^* : & 221\overline{144221}144 \\ \mathcal{C}_4^* : & 331\overline{122331}122 \end{array}$$

obtained from the family $\mathcal{C}$ by switching the blocks $22\overline{22}$ and $33\overline{33}$ in the cycle assigned to 1 and by leaving the other cycles invariant. To rule out $\mathcal{C}^*$ from the set of families of cycles of double pseudoline arrangements it remains to observe that the permutations that carry $\mathcal{C}_1$ onto $\mathcal{C}_1^*$ are exactly the 4 permutations $1324, 1243, 1432, \overline{1234}$ and that none of these 4 permutations leaves invariant the triplet $\mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_3$. Similarly if the chirotope of Γ is the chirotope of $M_2(12234)$ then the family of side cycles of Γ is either the family

$$\begin{array}{ll} \mathcal{C}_1 : & \overline{22}22\overline{33}33\overline{4}44\overline{4} \\ \mathcal{C}_2 : & \overline{33}1\overline{14}433\overline{1}14\overline{4} \\ \mathcal{C}_3 : & \overline{22}4\overline{4}1\overline{1}22\overline{4}4\overline{1}1 \\ \mathcal{C}_4 : & \overline{22}1\overline{1}2233\overline{1}1\overline{33} \end{array}$$

of side cycles of $M_2(1234)$ or the family

$$\begin{array}{ll} \mathcal{C}_1^* : & \overline{33}33\overline{22}22\overline{4}44\overline{4} \\ \mathcal{C}_2^* : & \overline{33}1\overline{14}433\overline{1}14\overline{4} \\ \mathcal{C}_3^* : & \overline{22}4\overline{4}1\overline{1}22\overline{4}4\overline{1}1 \\ \mathcal{C}_4^* : & \overline{22}1\overline{1}2233\overline{1}1\overline{33} \end{array}$$

obtained from the family $\mathcal{C}$ by switching the blocks $22\overline{22}$ and $33\overline{33}$ in the cycle assigned to 1 and by leaving the other cycles invariant. Again to rule out $\mathcal{C}^*$ from the set of families of cycles of double pseudoline arrangements it remains to observe that the permutations that carry $\mathcal{C}_1$ onto $\mathcal{C}_1^*$ are exactly the two permutations $1324, \overline{123}4$, and that none of these 2 permutations leaves invariant the cycle $\mathcal{C}_4$. $\square$

4.3. **Cocycles.** We repeat the definition of cocycles and introduce *cocycle-labels* as an alternative to the definition of cocycles.

Let Δ be a finite indexed family of pairwise disjoint oriented convex bodies of a projective plane $(\mathcal{P}, \mathcal{L})$, and let τ be a line of $(\mathcal{P}, \mathcal{L})$. Recall that we have defined

1. the *cocycle of* Δ *at* τ as the homeomorphism class of the image of the pair (Δ, τ) under the quotient map $\omega_\tau : \mathcal{P} \to \mathcal{P}/\mathcal{R}_\tau$ relative to the equivalence relation $\mathcal{R}_\tau$ whose nontrivial equivalence classes are the proper line segments of $\Delta \cap \tau$;
2. a *bitangent cocycle* or *zero-cocycle* as a cocycle at a bitangent;
3. the *isomorphism class of* Δ as the set of configurations Δ' that have the same set of bitangent cocycles as Δ; and the *chirotope of* Δ as the map that assigns to each triple J of the indexing set of Δ the isomorphism class of the subfamily indexed by J.

It is convenient to add to these three definitions a fourth one

4. the *cocycle map* of Δ as the map that assigns to each cell σ of the dual arrangement of Δ the cocycle of Δ at an element (hence any) of σ.

Figure 27 depicts, up to reorientation and reindexation of the convex bodies, the cocycles of families of two and three pairwise disjoint convex bodies with respective indexing set $\{1, 2\}$ and $\{1, 2, 3\}$; in this figure each circular diagram is labeled at its right bottom corner by its number of reoriented and reindexed versions and at its left bottom corner by a natural coding of the cocycle (composed of words on the alphabet obtained by adjoining to the indexing set of the configuration the negatives of its elements and the extra symbol $\bullet$; the negative of α is denoted $\overline{\alpha}$), called its *cocycle-label*, defined as follows. Let $\mathcal{D}_\tau$ be the closed 2-cell obtained by cutting $\mathcal{P}$ along the line τ, let $\nu_\tau : \mathcal{D}_\tau \to \mathcal{P}$ be the induced canonical projection, and let Σ_τ be the set of connected components of the preimages under ν_τ of the bodies of Δ; each element of Σ_τ is endowed with the orientation inherited from the orientation of the corresponding body in Δ. Choose an orientation ϵ of $\mathcal{D}_\tau$ and define the *cocycle-label* of the triple (Δ, τ, ϵ) as the set composed of

(1) the indices α whose corresponding convex bodies Δ_α are contained in the interior of $\mathcal{D}_\tau$ and oriented consistently with the orientation of $\mathcal{D}_\tau$;
(2) the negatives of the indices α whose corresponding convex bodies Δ_α are contained in the interior of $\mathcal{D}_\tau$ and oriented inconsistently with the orientation of $\mathcal{D}_\tau$; and
(3) the image of the circular sequence of elements of Σ_τ encountered when walking along the boundary of $\mathcal{D}_\tau$ according to its orientation under the morphism that assigns to a body its index or the negative of its index depending on whether the orientation of the body and the orientation of $\mathcal{D}_\tau$ are consistent or not, and to a point or line-segment the symbol $\bullet$.

The cocycle-label of the pair (Δ, τ) or the cocycle-label of Δ at τ is then defined as the cocycle-labels of the triples (Δ, τ, ϵ) and $(\Delta, \tau, -\epsilon)$, and can be represented by any of its two elements since the cocycle-label of the triple $(\Delta, \tau, -\epsilon)$ is obtained from the cocycle-label of the triple (Δ, τ, ϵ) by replacing each of its element by the reversal of its overlined version (with the usual convention that an index coincides with the negative of its negative : $\overline{\overline{\alpha}} = \alpha$). Clearly the cocycle of a pair (Δ, τ) depends only on its cocycle-label and vice-versa. Figure 28 depicts examples of cocycle maps of families of one, two, and three pairwise disjoint convex bodies with respective indexing sets $\{1\}$, $\{1, 2\}$, and $\{1, 2, 3\}$. In particular one can easily check that the cocycle map of a family of two bodies is one-to-one.

Lemma 24. *Cocycle maps of families of two disjoint convex bodies are one-to-one.* □

We are now ready to prove that the duality map is compatible with the isomorphism relations on families of convex bodies and on arrangements of double pseudolines, and induces a one-to-one (and onto) correspondence between their quotient sets. This means, for example, that the bitangent cocycles

$$\{1\bullet\overline{2}\bullet\overline{3}\bullet\}, \{1\bullet\overline{3}\bullet2\bullet\}, \{1\bullet2\bullet3\bullet\}, \{1\bullet3\bullet\overline{2}\bullet\}$$

of the configuration of three convex bodies depicted at the left bottom part of Figure 28 is a coding of the isomorphism class of the dual arrangement of the configuration.



$12\overline{12}$ 1
$1\overline{1}, 2$ 2
$12\bullet\bullet$ 4
$\overline{2}12\bullet$ 4
$1\bullet, 2$ 4
$1, 2$ 2
$1\bullet2\bullet3\bullet$ 8
$123\bullet\bullet\bullet$ 24
$1\overline{3}2\bullet3\bullet$ 24
$12\bullet\bullet, 3$ 24
$\overline{3}123\bullet\bullet$ 24
$1\bullet, 2, 3$ 12
$\overline{2}12\bullet, 3$ 12
$\overline{23}123\bullet$ 24
$1, 2, 3$ 4
$1\overline{1}, 2, 3$ 8
$12\overline{12}, 3$ 6
$123\overline{123}$ 4

FIGURE 27. Cocycles of indexed families of two and three pairwise-disjoint oriented convex bodies.



FIGURE 28. Indexed families of one, two and three pairwise-disjoint oriented convex bodies and the cocycle labeled versions of their dual arrangements.

Theorem 25. *Let Δ and Δ' be two finite families of pairwise disjoint oriented convex bodies of projective planes. Then the four following assertions are equivalent*

(1) Δ and Δ' have the same chirotope;
(2) Δ and Δ' have isomorphic dual arrangements;
(3) Δ and Δ' have isomorphic cocycle maps;
(4) Δ and Δ' have the same set of 0-cocycles (i.e., are isomorphic). □

Proof. Some implications are clear:

(i) $(4) \Rightarrow (1)$;
(ii) $(4), (2) \Rightarrow (3)$, using a perturbation argument;
(iii) $(2), (2) \Rightarrow (4)$, because the family of 0-cocycles of Δ depends only the family of cocycle-labeled versions of the dual arrangements of subfamilies of three bodies and on the isomorphism class of the dual arrangement of Δ;
(iv) $(3) \Rightarrow (4), (2), (1)$.

We now prove that $(1) \Leftrightarrow (2)$.

We first prove that $(2) \Rightarrow (1)$. Let $\mathcal{V}$ be the (finite) set of cocycle-labels $\mu(\Delta, \tau)$ of the pairs (Δ, τ) as Δ ranges over the set of families of $n \geq 3$ convex bodies indexed by $\{1, 2, 3, \ldots, n\}$ and where τ ranges over the set of bitangents of Δ. We leave the verification of the following property of the set $\mathcal{V}$ to the reader: the map that assigns to any element $\mu(\Delta, \tau)$ of $\mathcal{V}$ the set of $\mu(\Delta', \tau)$ where Δ' ranges over the set of subfamilies of size $n-1$ of Δ is one-to-one: see Table 2 for the case $n = 3$; this proves that $(2) \Rightarrow (1)$.

$\mu(123, \tau)$	$\mu(12, \tau)$	$\mu(13, \tau)$	$\mu(23, \tau)$
$123\bullet\bullet\bullet$	$12\bullet\bullet$	$13\bullet\bullet$	$23\bullet\bullet$
$1\bullet2\bullet3\bullet$	$12\bullet\bullet$	$31\bullet\bullet$	$23\bullet\bullet$
$12\bullet\bullet, 3$	$12\bullet\bullet$	$1\bullet, 3$	$2\bullet, 3$
$132\bullet\overline{3}\bullet$	$12\bullet\bullet$	$\overline{3}13\bullet$	$32\overline{3}\bullet$
$\overline{3}123\bullet\bullet$	$12\bullet\bullet$	$\overline{3}13\bullet$	$\overline{3}23\bullet$

Table 2. The map that assigns to a 0-cocycle of a family of three convex bodies its sub-cocycles on two bodies is one-to-one.

We now prove that $(1) \Rightarrow (2)$. It is sufficient to prove it for families of three bodies. Let I be the indexing set of Δ, let $\mathcal{I}$ be the set of pairs (i, J) where i ranges over I and where J ranges over the set of 3-subsets of I that contains i, and for $(i, J) \in \mathcal{I}$ let $C_{i,J}$ be the circular ordering of the bitangents v_α, $\alpha \in \{ij, \overline{i}j, i\overline{j}, \overline{ij} \mid j \in I \setminus i\}$, along the (oriented) dual curve Δ_i^* of Δ_i. According to Theorem 15 the isomorphism class of the dual arrangement of Δ depends only on the family of $C_{i,J}$, $(i, J) \in \mathcal{I}$. Thus proving that $(1) \Rightarrow (2)$ comes down to proving that the $C_{i,J}$ depend only on the chirotope of Δ. This latter statement is a simple consequence of the two following observations

(1) for any $j \in I \setminus \{i\}$, the four vertices v_α, $\alpha \in \{ij, \overline{i}j, i\overline{j}, \overline{ij}\}$, appear by definition in the circular order v_{ij},$v_{\overline{i}j}$,$v_{i\overline{j}}$,$v_{\overline{ij}}$ along Δ_i^*;

(2) for any $j \in I \setminus \{i\}$ and any $\alpha \in \{ij, \bar{i}j, i\bar{j}, \overline{ij}\}$ the position of v_α with respect to the v_β, $\beta \in \{ik, \bar{i}k, i\bar{k}, \overline{ik}\}$, $k \in I \setminus \{i, j\}$, depends only on the chirotope of Δ for the cocycle map is one-to-one for families of two bodies.

□

Figures 29, 30 and 31 depict the zero-cocycle labeled versions of the thirteen indexed and oriented simple arrangements on three double pseudolines of Figures 25 and 26.


$123\bullet\bar{2}\bullet$
$3\bar{2}1\bullet2\bullet$
$3\overline{12}\bullet1\bullet$
$2\bar{1}3\bullet1\bullet$
2
$\overline{23}1\bullet3\bullet$
3
$1\bar{3}2\bullet3\bullet$
1
$13\bar{2}\bullet\bar{3}\bullet$
$231\bullet\bar{3}\bullet$
$\overline{213}\bullet1\bullet$
$\overline{312}\bullet1\bullet$
C_{64}
$\bar{3}21\bullet\bar{2}\bullet$
$1\overline{23}\bullet2\bullet$
2

FIGURE 29. The cocycle labelled version of the arrangement $C_{64}(123)$.


$\overline{2}1\bullet\bullet, 3$
$12\bullet\bullet, 3$
$3\overline{2}\bullet\bullet, 1$
$23\bullet\bullet, 1$
$13\bullet\bullet, \overline{2}$
$31\bullet\bullet, 2$
1
3
2
$\overline{3}1\bullet\bullet, \overline{2}$
$1\overline{3}\bullet\bullet, 2$
$\overline{23}\bullet\bullet, 1$
049000
$\overline{3}2\bullet\bullet, 1$
$1\overline{2}\bullet\bullet, \overline{3}$
$21\bullet\bullet, \overline{3}$
$C_{04}(123)$
2
$\overline{2}1\bullet\bullet, 3$
$12\bullet\bullet, 3$
$3\overline{2}\bullet\bullet, 1$
$23\bullet\bullet, 1$
$13\bullet\bullet, \overline{2}$
$31\bullet\bullet, 2$
1
3
2
$\overline{3}1\bullet\bullet, \overline{2}$
$231\bullet\overline{3}\bullet$
$\overline{23}\bullet\bullet, 1$
073300
$\overline{31}2\bullet1\bullet$
$1\overline{2}\bullet\bullet, \overline{3}$
$1\overline{23}\bullet2\bullet$
C_{07}
8
$\overline{2}1\bullet\bullet, 3$
$3\overline{2}1\bullet2\bullet$
$3\overline{2}\bullet\bullet, 1$
$2\overline{1}3\bullet1\bullet$
$13\bullet\bullet, \overline{2}$
$1\overline{3}2\bullet3\bullet$
2
3
1
$\overline{3}1\bullet\bullet, \overline{2}$
$231\bullet\overline{3}\bullet$
$\overline{23}\bullet\bullet, 1$
181030
$\overline{31}2\bullet1\bullet$
$1\overline{2}\bullet\bullet, \overline{3}$
$1\overline{23}\bullet2\bullet$
C_{18}
12
$\overline{2}1\bullet\bullet, 3$
$3\overline{2}1\bullet2\bullet$
$3\overline{2}\bullet\bullet, 1$
$2\overline{1}3\bullet1\bullet$
$13\bullet\bullet, \overline{2}$
$1\overline{3}2\bullet3\bullet$
1
2
3
$13\overline{2}\bullet\overline{3}\bullet$
$231\bullet\overline{3}\bullet$
$\overline{213}\bullet1\bullet$
370003
$\overline{31}2\bullet1\bullet$
$\overline{3}21\bullet\overline{2}\bullet$
$1\overline{23}\bullet2\bullet$
C_{37}
8
$31\bullet\bullet, \overline{2}$
$3\overline{2}1\overline{3}\bullet\bullet$
$\overline{1}3\overline{2}1\bullet\bullet$
$2\overline{1}3\bullet1\bullet$
$13\bullet\bullet, \overline{2}$
$1\overline{3}2\bullet3\bullet$
2
3
1
$\overline{3}1\bullet\bullet, \overline{2}$
$1\overline{3}\bullet\bullet, 2$
$\overline{23}\bullet\bullet, 1$
154300
$\overline{3}2\bullet\bullet, 1$
$1\overline{2}\bullet\bullet, \overline{3}$
$21\bullet\bullet, \overline{3}$
C_{15}
24
$31\bullet\bullet, \overline{2}$
$3\overline{2}1\overline{3}\bullet\bullet$
$\overline{1}3\overline{2}1\bullet\bullet$
$2\overline{1}3\bullet1\bullet$
$13\bullet\bullet, \overline{2}$
$1\overline{3}2\bullet3\bullet$
2
3
1
$13\overline{2}\bullet\overline{3}\bullet$
$231\bullet\overline{3}\bullet$
$\overline{213}\bullet1\bullet$
433021
$\overline{31}2\bullet1\bullet$
$\overline{3}21\bullet\overline{2}\bullet$
$1\overline{23}\bullet2\bullet$
C_{43}
24

FIGURE 30. Cocycle labelled versions of the arrangements $C_{04}, C_{07}, C_{18}, C_{37}, C_{15}, C_{43}$.


$31\bullet\bullet, \overline{2}$
$1\overline{3}\bullet\bullet, \overline{2}$
$\overline{1}3\overline{2}1\bullet\bullet$
$2\overline{1}3\bullet 1\bullet$
$13\bullet\bullet, \overline{2}$
$\overline{2}1\bullet\bullet\overline{3}$
2
3
1
$\overline{3}1\bullet\bullet, \overline{2}$
$12\bullet\bullet, \overline{3}$
$\overline{2}1\overline{3}\bullet\overline{1}\bullet$
22_{8010}
$1\overline{3}2\overline{1}\bullet\bullet$
$1\overline{2}\bullet\bullet, \overline{3}$
$21\bullet\bullet, \overline{3}$
C_{22}
12
$31\bullet\bullet, \overline{2}$
$3\overline{2}1\overline{3}\bullet\bullet$
$\overline{1}3\overline{2}1\bullet\bullet$
$2\overline{1}3\bullet 1\bullet$
$13\bullet\bullet, \overline{2}$
$\overline{2}1\overline{3}2\bullet\bullet$
2
3
1
$13\overline{2}\bullet\overline{3}\bullet$
$12\bullet\bullet, \overline{3}$
$\overline{213}\bullet 1\bullet$
33_{3310}
$1\overline{3}2\overline{1}\bullet\bullet$
$\overline{3}21\bullet\overline{2}\bullet$
$21\bullet\bullet, \overline{3}$
C_{33}
24
$31\bullet\bullet, \overline{2}$
$3\overline{2}1\overline{3}\bullet\bullet$
$\overline{1}3\overline{2}1\bullet\bullet$
$2\overline{1}3\bullet 1\bullet$
$13\bullet\bullet, \overline{2}$
$1\overline{3}2\bullet 3\bullet$
2
3
1
$\overline{3}1\bullet\bullet, \overline{2}$
$231\bullet\overline{3}\bullet$
$1\overline{231}\bullet\bullet$
32_{6020}
$\overline{31}2\bullet 1\bullet$
$1\overline{3}\bullet\bullet, \overline{2}$
$31\overline{23}\bullet\bullet$
C_{32}
24
$31\bullet\bullet, \overline{2}$
$3\overline{2}1\overline{3}\bullet\bullet$
$\overline{1}3\overline{2}1\bullet\bullet$
$2\overline{1}3\bullet 1\bullet$
$13\bullet\bullet, \overline{2}$
$\overline{2}1\overline{3}2\bullet\bullet$
2
3
1
$\overline{3}1\bullet\bullet, \overline{2}$
$12\bullet\bullet, \overline{3}$
$\overline{23}\bullet\bullet, 1$
25_{2310}
2
$1\overline{3}2\overline{1}\bullet\bullet$
$1\overline{2}\bullet\bullet, \overline{3}$
$21\bullet\bullet, \overline{3}$
C_{25_2}
24
$\overline{2}1\bullet\bullet, 3$
$3\overline{2}1\bullet 2\bullet$
$3\overline{2}\bullet\bullet, 1$
$2\overline{1}3\bullet 1\bullet$
$13\bullet\bullet, \overline{2}$
$\overline{2}1\overline{3}2\bullet\bullet$
1
3
2
$13\overline{2}\bullet\overline{3}\bullet$
$12\bullet\bullet, \overline{3}$
$\overline{213}\bullet 1\bullet$
25_{2310}
1
$1\overline{3}2\overline{1}\bullet\bullet$
$\overline{3}21\bullet\overline{2}\bullet$
$21\bullet\bullet, \overline{3}$
C_{25_1}
48
$31\bullet\bullet, \overline{2}$
$213\overline{2}\bullet\bullet$
$\overline{1}3\overline{2}1\bullet\bullet$
$23\bullet\bullet, 1$
2
$13\bullet\bullet, \overline{2}$
3
$\overline{2}132\bullet\bullet$
$\overline{3}1\bullet\bullet, \overline{2}$
1
$12\bullet\bullet, \overline{3}$
$\overline{23}\bullet\bullet, 1$
36_{0040}
$1\overline{3}2\overline{1}\bullet\bullet$
$21\bullet\bullet, \overline{3}$
$1\overline{2}\bullet\bullet, \overline{3}$
C_{36}
4

FIGURE 31. Cocycle labelled versions of arrangements $C_{22}, C_{33}, C_{32}, C_{25_2}, C_{25_1}$ and C_{36}.

5. AXIOMATIC CHARACTERIZATION

In this section we give a proof of the axiomatic characterization of the class of chirotopes of double pseudoline arrangements announced in the introduction; cf. Theorem 4.

Let $k \geq 3$ be a natural number. It is convenient to use the following terminology. A *chirotope* is a map χ that assigns to each triple of a finite indexing set I of at least three indices an isomorphism class of double pseudoline arrangements indexed by that triple (an entry of the chirotope for short); a *k-extension* of a chirotope χ, defined on the set of triples of the indexing set I, is the extension of χ (if any) to the set of subsets J of I of size k that assigns to J the isomorphism class of the arrangements of double pseudolines whose chirotopes are the restriction of χ to the triples of J; and a *k-chirotope* is a chirotope which admits a k-extension, i.e., whose restrictions to the sets of triples of subsets of indices of size 3 to k are chirotopes of double pseudoline arrangements. For example the chirotope χ on the indexing set $\{1, 2, 3, 4, 5\}$ with entries

$$\begin{matrix} C_{04}(123) & C_{04}(124) & C_{04}(125) & C_{04}(134) & C_{04}(145) \\ C_{04}(234) & C_{04}(245) & C_{04}(345) & C_{04}(153) & C_{04}(253) \end{matrix}$$

admits a 4-extension, depicted in Figure 32, but no 5-extension because the side cycles



1234 1245 1235

1345 2345

FIGURE 32. The 4-extension of a 4-chirotope on the indexing set $1, 2, 3, 4, 5$.

of disk type $23\overline{23}, 24\overline{24}, 25\overline{25}, 34\overline{34}, 45\overline{45}$, $53\overline{53}$ assigned to the index 1 of the entries $C_{04}(123)$, $C_{04}(124)$, $C_{04}(125)$, $C_{04}(134)$, $C_{04}(145)$ and $C_{04}(153)$ of χ are not *compatible*: not compatible in the sense that there is no cycle involving the indices $2, 3, 4, 5$ and their negatives exactly once in which $23\overline{23}, 24\overline{24}, 25\overline{25}, 34\overline{34}, 45\overline{45}$ and $53\overline{53}$ would be subcycles.

Thus our axiomatization theorem simply states that the class of 5-chirotopes is the class of chirotopes of double pseudoline arrangements. To prove our axiomatization theorem we define a class of operators on the set of 5-chirotopes, compatible with the property "to be a chirotope of double pseudoline arrangements" and we show that any 5-chirotope is the image of the chirotope of the double version of a cyclic arrangement of pseudolines under a finite product of such operators. These operators are defined as conjugates of mutation operators (i.e., inversion of a triangular face and, more generally, inversion of a fan) on a certain class of finite arrangements of simple closed curves embedded in compact nonorientable surfaces in one-to-one correspondence with the class of 5-chirotopes.

5.1. **Arrangements of genus** $1, 2, \ldots$ **.** We extend the definition of arrangements of double pseudolines by relaxing the condition on the arrangement which says that the genus of its underlying surface is 1. By an indexed and oriented arrangement (of double pseudolines) of genus $g \geq 1$ we mean a finite indexed family τ of at least two simple closed oriented curves embedded in a compact nonorientable surface S_τ of genus g with the property that there exists closed tubular neighborhoods R_i of the τ_i (ribbons for short) such that

(1) for any subfamily ν of τ the union of its ribbons, denoted, R_ν, is a closed tubular neighborhood of the union of its curves; the compact surface obtained by attaching, using homeomorphisms for the attaching maps, topological disks to the boundary curves of R_ν is denoted S_ν;
(2) any subfamily ν of τ of size 2 considered as embedded not in S_τ but in S_ν is homeomorphic to the dual arrangement of a (hence any) configuration of two convex bodies;
(3) for any $\tau_i, \tau_j \in \tau$ the intersection of the disk side of τ_i in the subarrangement τ_i, τ_j and the ribbon R_i is independent of τ_j.

Thus arrangements of genus 1 are the arrangements of double pseudolines as defined in the previous sections. Figure 33 depicts two embeddings in 3-space of the tubular neighborhood of an arrangement of two double pseudolines (thus a union of two ribbons). A horizontal dashed line segment indicates the presence of a half-twist (180 degree) of



Figure 33.

the ribbon crossed by the line segment and the numbers, in the right diagram, label the corners of the polygonal boundary curves of the neighborhood: the corners of a polygonal boundary curve being labeled by the same number. We extend in the natural way to the class of arrangements of arbitrary genus the notions of thinness, mutations, isomorphism classes, node cycles, side cycles of disk and crosscap type, and chirotopes associated with the class of arrangements of genus 1. As for arrangements of genus 1 the isomorphism class of an arrangement of genus g depends only on its family of side cycles, with the net benefit that there is now a simple characterization of cycles that arose as side cycles of arrangements. Given a sequence $B = \{i_1j_1\}\{i_2j_2\}\{i_3j_3\}\dots\{i_kj_k\}$, $i_l \in \pm i$, we denote by $B_{\otimes j_p}$ the sequence $\alpha_{p+1}\alpha_{p+2}\dots\alpha_{k_m}\alpha_p\alpha_1\alpha_2\dots\alpha_{p-1}$ or its reverse $\alpha_{p-1}\alpha_{p-2}\dots\alpha_2\alpha_1\alpha_p\alpha_{k_m}\dots\alpha_{p+2}\alpha_{p+1}$, depending on whether $i_p = \bar{i}$ or $i_p = i$, where

$$\alpha_q = \begin{cases} \{i_pj_{mp}\} \otimes \{i_qj_{mq}\} & \text{if } p < q \leq k_m \\ \{i_pj_{mp}\} & \text{if } q = p \\ \{i_qj_{mq}\} \otimes \{i_pj_{mp}\} & \text{if } 1 \leq q < p. \end{cases}$$

We leave the verification of the following property to the reader: if B is a prime factor of the side cycle of disk type indexed by i of an arrangement then $B_{\otimes j_p}$ is the corresponding prime factor of the side cycle of disk type indexed by j_{mp}. Routine considerations that we leave to the reader result in the following theorem.

Theorem 26. *Let I be a finite set of (at least 2) indices, let D_i and M_i, $i \in I$, be two families of circular words on the signed version $\hat{I}$ of I, with the property that D_i and M_i are shuffles of the elementary cycles $jj\overline{jj}$, $j \neq i$, let S_i be the result of replacing in D_i the linear subsequences $\overline{jj}jj$, $j \neq i$, by the linear sequences $\{ij\}\{\bar{i}j\}\{i\bar{j}\}\{\overline{ij}\}$, let T_i be the result of replacing in M_i linear subsequences $\overline{jj}jj$, $j \neq i$, by the linear sequences $\{ij\}\{\bar{i}j\}\{i\bar{j}\}\{\overline{ij}\}$, and let $S_{\bar{i}}$ and $T_{\bar{i}}$ be the reversal of S_i and T_i, respectively. Then the D_i and M_i are the side cycles of disk type and crosscap type of an arrangement Γ of oriented double pseudolines indexed by I if and only if there exists block decompositions*

$$B_{i1}B_{i2}\dots B_{in_i}$$

of the S_i, $i \in I$, where $B_{im} = \{i_1j_{m1}\}\{i_2j_{m2}\}\{i_3j_{m3}\}\dots\{i_kj_{mk_m}\}$, $1 \leq m \leq n_i$, with $j_{m_l} \neq \pm j_{m_{l'}}$ for all $1 \leq l < l' \leq k_m$, such that

1. *$T_i = B^*_{i1}B^*_{i2}\dots B^*_{in_i}$ where B^*_{im} is the reversal of B_{im} (note that the pair S_i, T_i determine their block decompositions);*
2. *$B_{\otimes j_p}$ is one of the blocks of the block decomposition of $S_{j_{mp}}$.* ☐

An arrangement τ is called a *k-arrangement* if any subfamily ν of τ of size k is an arrangement of genus one. Clearly chirotopes of k-arrangements are k-chirotopes. Before proving that the converse is true for $k \geq 5$ we give few examples.

Example 7. Figure 34 depicts a family of three curves embedded in a Klein bottle (decomposed by the curves into 2 digons, 2 trigons, 6 tetragons, 1 hexagon and 1 octagon) that fulfills condition (2) but not condition (3) of the definition of an arrangement : the disk side of the green curve in the arrangement composed of the green and red curve and the disk side of the green curve in the arrangement composed of the green and black curve intersect the ribbon of the green curve into two distinct cuffs. (On the other hand disk and crosscap sides of the red and black curves are well-defined.)



FIGURE 34.

Example 8. Figure 35 depicts an arrangement of three curves, obtained by adding two twists on the green curve of the configuration of Figure 34. It is composed of 2 digons, 6 tetragons, 1 octagon and 1 dodecagon (the corners of the octagon are labeled with the numeral 8 and those of the dodecagon by the letter γ). It lives in a double Klein bottle. Its node cycles are the same as those of the previous example, i.e.,



FIGURE 35.

$$\begin{array}{rl}1: & 12, \overline{1}2, 1\overline{2}, \overline{12}, 1\overline{3}, \overline{13}, 13, \overline{1}3\\ 2: & 21, \overline{2}1, 2\overline{1}, \overline{21}, 2\overline{3}, \overline{23}, 23, \overline{2}3\\ 3: & 3\overline{1}, 3\overline{2}, \overline{32}, \overline{31}, 31, 32, \overline{3}2, \overline{3}1\end{array}$$

Example 9. Figure 36 depicts embeddings in 3-space of tubular neighborhoods of two arrangements on three curves. Again the horizontal dashed line segments indicate the presence of half-twists (180 degree) of the ribbons of the tubular neighborhood. Both



FIGURE 36.

live in a sphere with 4 crosscaps (a double Klein bottle) decomposed by the curves into 1 trigon, 7 tetragons, 1 octagon and 1 nonagon. (In the left diagram the corners of the octagon are labeled by the numeral 8 and those of the nonagon by the numeral 9.) If we orient clockwise the curves and use, respectively, the indices 1, 2 and 3 for the green, blue and red curves, then the side cycles of disk type of the arrangements are, respectively,

$$\begin{array}{ll} 1: & \overline{22}22\overline{33}33 \\ 2: & \overline{3311}3311 \\ 3: & \overline{22}1122\overline{11} \end{array}$$

and

$$\begin{array}{ll} 1: & \overline{22}3322\overline{33} \\ 2: & \overline{331}1331\overline{1} \\ 3: & \overline{221}1221\overline{1} \end{array}$$

Observe that the first arrangement is a martagon (with respect to and only to the green curve) but not the second one and that these two arrangements are connected by a sequence of four mutations.

Example 10. Figure 37 depicts the cell complex of a simple arrangement on three octagonal curves (colored red, green and purple in pdf color)

$$\begin{array}{rcl} \tau_1 & = & 1a2b3c4d5e6f7g8h \\ \tau_2 & = & \hat{1}\hat{a}\hat{2}\hat{b}\hat{3}\hat{c}\hat{4}\hat{d}\hat{5}\hat{e}\hat{6}\hat{f}\hat{7}\hat{g}\hat{8}\hat{h}, \\ \tau_3 & = & \tilde{1}\tilde{a}\tilde{2}\tilde{b}\tilde{3}\tilde{c}\tilde{4}\tilde{d}\tilde{5}\tilde{e}\tilde{6}\tilde{f}\tilde{7}\tilde{g}\tilde{8}\tilde{h}, \end{array}$$

living in a triple cross surface as one can check by calculating the Euler characteristic of the surface. In the Figure the cell complex is augmented with its dual graph (oriented arbitrarily at our convenience): Using the symbol of an edge of the cell complex to denote its dual we get a dual presentation composed of a system of 12 equations in 24 symbols

$$\begin{array}{llll} a\hat{f} = \hat{e}b & a\hat{h} = \hat{g}h & c\hat{f} = \hat{g}b & c\hat{d} = \hat{e}d \\ \tilde{a}f = e\tilde{b} & \tilde{a}h = g\tilde{h} & \tilde{c}f = g\tilde{b} & \tilde{c}d = e\tilde{d} \\ \hat{a}\tilde{f} = \tilde{e}\hat{b} & \hat{a}\tilde{h} = \tilde{g}\hat{h} & \hat{c}\tilde{f} = \tilde{g}\hat{b} & \hat{c}\tilde{d} = \tilde{e}\hat{d} \end{array}$$



FIGURE 37. A 2-arrangement of three double pseudolines living in a triple cross surface. The double pseudolines are drawn red, green and purple in pdf color.

putting in evidence that this system of curves is well-defined arrangement, as illustrated in Figure 38 where the shaded regions stand for the crosscap sides of the curves. We built it as the simple arrangement with side cycles (of disk type)

$$\begin{array}{ll} 1: & 22\overline{22}33\overline{33} \\ 2: & 33\overline{33}11\overline{11} \\ 3: & 11\overline{11}22\overline{22} \end{array}$$

(this can be read easily on the dual presentation). Observe that it is a martagon with respect to each of its three curves.

The number of simple arrangements on three curves on a given indexing set is 140^3, the cube of the number $4\binom{7}{3}$ of possible cycles of length 8 (i.e., the number of shuffles of the



FIGURE 38. A 2-arrangement of three double pseudolines living in a triple cross surface.

circular sequences $\overline{22}22$ and $\overline{33}33$). More generally the number of simple arrangements on n curves is the n-th power of the number

$$\begin{aligned} s_n &= \frac{1}{(4n-4)}\binom{4n-4}{4}\binom{4n-8}{4}\cdots\binom{8}{4}\binom{4}{4} \\ &= \frac{1}{(4n-4)}\frac{(4n-4)!}{(4!)^{n-1}} \end{aligned}$$

of shuffles of the $n-1$ circular sequences $\overline{jj}jj$, $1 \leq j \leq n$, $j \neq 1$.

Example 11. The thin 4-chirotope on five indices depicted in Figure 32 is not a 4-arrangement: actually, the only thin 4-arrangement on five indices is the cyclic arrangement on five double pseudolines.

Example 12. The simple 3-chirotope on the 4-set of indices $\{1,2,3,4\}$ with entries $C_{64}(123), C_{64}(124), C_{64}(134), C_{64}(234)$ is the chirotope of a unique simple arrangement Υ on four curves whose side cycles of disk type are

$$\begin{aligned} 1:&\quad 2\overline{2}3\overline{3}4\overline{42}2\overline{3}3\overline{4}4 \\ 2:&\quad 3\overline{3}4\overline{4}1\overline{13}3\overline{4}4\overline{1}1 \\ 3:&\quad 4\overline{4}1\overline{1}2\overline{24}4\overline{1}1\overline{2}2 \\ 4:&\quad 1\overline{1}2\overline{2}3\overline{31}1\overline{2}2\overline{3}3. \end{aligned}$$

The surface is a sphere with 7 crosscaps decomposed by the curves into 19 two-cells (12 digons, 3 octagons, and 4 dodecagons) put together according to the following presentation

$$\begin{array}{rrlrrl}
1: & a\hat{a}^{-1} & =1 & 2: & g\hat{g} & =1 \\
3: & \hat{i}\tilde{f}^{-1} & =1 & 4: & \hat{c}\tilde{l}^{-1} & =1 \\
5: & \tilde{b}\check{f}^{-1} & =1 & 6: & \tilde{h}\check{l} & =1 \\
7: & e\check{b}^{-1} & =1 & 8: & k\check{h} & =1 \\
9: & c\tilde{d}^{-1} & =1 & 10: & i\tilde{j} & =1 \\
11: & \hat{k}\check{d}^{-1} & =1 & 12: & \hat{e}\check{j} & =1 \\
13: & \hat{b}^{-1}b\tilde{c}^{-1}\check{g}l\hat{l}^{-1}\check{e}\tilde{a}^{-1} & =1 & & & \\
14: & \check{a}\tilde{g}\hat{j}^{-1}\check{c}f^{-1}\hat{h}^{-1}\tilde{e}d^{-1} & =1 & 15: & \hat{f}h\tilde{k}\hat{d}\check{k}\tilde{i}j\check{i} & =1 \\
16: & a\hat{b}\tilde{l}^{-1}\hat{d}\check{j}^{-1}\hat{f}g^{-1}\hat{h}\tilde{f}\hat{j}\check{d}\hat{l} & =1 & 17: & \hat{a}b\tilde{d}d\check{b}f\hat{g}^{-1}h\tilde{j}^{-1}j\check{h}^{-1}l & =1 \\
18: & \tilde{e}\hat{i}\tilde{g}\check{l}^{-1}\tilde{i}i^{-1}\tilde{k}\hat{c}^{-1}\tilde{a}\check{f}\tilde{c}c & =1 & 19: & \hat{c}\hat{k}\check{e}\tilde{b}\check{g}k^{-1}\check{i}\hat{e}^{-1}\check{k}\tilde{h}^{-1}\check{a}e & =1
\end{array}$$

where

$$\begin{array}{rcl}
\Upsilon_1 & = & abcdefghijkl \\
\Upsilon_2 & = & \hat{a}\hat{b}\hat{c}\hat{d}\hat{e}\hat{f}\hat{g}\hat{h}\hat{i}\hat{j}\hat{k}\hat{l} \\
\Upsilon_3 & = & \tilde{a}\tilde{b}\tilde{c}\tilde{d}\tilde{e}\tilde{f}\tilde{g}\tilde{h}\tilde{i}\tilde{j}\tilde{k}\tilde{l} \\
\Upsilon_4 & = & \check{a}\check{b}\check{c}\check{d}\check{e}\check{f}\check{g}\check{h}\check{i}\check{j}\check{k}\check{l}.
\end{array}$$

A dual presentation is given by the following system of 24 equations in 48 symbols

$$\begin{array}{cccc}
a\hat{b}=\hat{a}b & \hat{g}f=g\hat{h} & g\hat{f}=\hat{g}f & a\hat{l}=\hat{a}l \\
c\tilde{e}=\tilde{d}d & \tilde{j}h=i\tilde{k} & i\tilde{i}=\tilde{j}j & \tilde{d}b=c\tilde{c} \\
e\check{c}=\check{b}f & \check{h}j=k\check{i} & k\check{g}=\check{h}l & \check{b}d=e\check{a} \\
\hat{i}\tilde{g}=\tilde{f}\hat{j} & \hat{c}\tilde{a}=\tilde{l}\hat{b} & \hat{c}\tilde{k}=\tilde{l}\hat{d} & \tilde{f}\hat{h}=\hat{i}\tilde{e} \\
\hat{k}\check{e}=\check{d}\hat{l} & \check{j}\hat{d}=\hat{e}\check{k} & \hat{e}\check{i}=\check{j}\hat{f} & \check{d}\hat{j}=\hat{k}\check{c} \\
\tilde{b}\check{g}=\check{f}\tilde{c} & \check{l}\tilde{g}=\tilde{h}\check{a} & \tilde{h}\check{k}=\check{l}\tilde{i} & \check{f}\tilde{a}=\tilde{b}\check{e}
\end{array}$$

where we use the same symbol to denote an edge and its dual; the dual presentation is also depicted in Figure 39.


1 $\hat{a}$ 17 a b 16 $\hat{b}$ 13
17 $\hat{g}$ 2 f g 14 $\hat{h}$ 16
16 $\hat{f}$ 15 g h 2 $\hat{g}$ 17
13 $\hat{l}$ 16 l a 17 $\hat{a}$ 1
9 $\tilde{d}$ 17 c d 18 $\tilde{e}$ 14
17 $\tilde{j}$ 10 h i 15 $\tilde{k}$ 18
18 $\tilde{i}$ 15 i j 10 $\tilde{j}$ 17
13 $\tilde{c}$ 18 b c 17 $\tilde{d}$ 9
7 $\check{b}$ 17 e f 19 $\check{c}$ 14
17 $\check{h}$ 8 j k 15 $\check{i}$ 19
19 $\check{g}$ 13 k l 8 $\check{h}$ 17
14 $\check{a}$ 19 d e 17 $\check{b}$ 7
3 $\tilde{f}$ 16 $\hat{i}$ $\hat{j}$ 18 $\tilde{g}$ 14
16 $\tilde{l}$ 4 $\hat{b}$ $\hat{c}$ 13 $\tilde{a}$ 18
18 $\tilde{k}$ 15 $\hat{c}$ $\hat{d}$ 4 $\tilde{l}$ 16
14 $\tilde{e}$ 18 $\hat{h}$ $\hat{i}$ 16 $\tilde{f}$ 3
11 $\check{d}$ 16 $\hat{k}$ $\hat{l}$ 19 $\check{e}$ 13
16 $\check{j}$ 12 $\hat{d}$ $\hat{e}$ 15 $\check{k}$ 19
19 $\check{i}$ 15 $\hat{e}$ $\hat{f}$ 12 $\check{j}$ 16
14 $\check{c}$ 19 $\hat{j}$ $\hat{k}$ 16 $\check{d}$ 11
5 $\check{f}$ 18 $\tilde{b}$ $\tilde{c}$ 19 $\check{g}$ 13
18 $\check{l}$ 6 $\tilde{g}$ $\tilde{h}$ 14 $\check{a}$ 19
19 $\check{k}$ 15 $\tilde{h}$ $\tilde{i}$ 6 $\check{l}$ 18
13 $\check{e}$ 19 $\tilde{a}$ $\tilde{b}$ 18 $\check{f}$ 5

FIGURE 39. $4 \times C_{64}$

Example 13. The chirotope of the martagon $M_1(1234)$ is the chirotope of a second martagon $M_1^*(1234)$ defined by the (side) cycles (of disk type)

$$\begin{array}{rl} 1: & \overline{22}22\overline{33}33\overline{44}44 \\ 2: & \overline{3344}1\overline{1}3344\overline{1}1 \\ 3: & \overline{4422}1\overline{1}4422\overline{1}1 \\ 4: & \overline{2233}1\overline{1}2233\overline{1}1 \end{array}$$

which are obtained from the cycles of $M_1(1234)$ by simply changing the order of the blocks $\overline{2}222$, $\overline{3}333$, and $\overline{4}444$ in the cycle indexed by 1. This arrangement lives in a triple cross surface that is decomposed by the curves into 3 digons, 15 tetragons, 3 pentagons, 1 hexagon and 1 nonagon. Note that this arrangement has no triangular faces. Similarly the chirotope of the martagon $M_2(1234)$ is the chirotope of a second martagon $M_2^*(1234)$ defined by the cycles

$$\begin{array}{rl} 1: & \overline{22}22\overline{4}44\overline{433}33 \\ 2: & \overline{33}1\overline{14}433\overline{1}14\overline{4} \\ 3: & \overline{22}4\overline{4}1\overline{1}22\overline{4}4\overline{1}1 \\ 4: & \overline{22}1\overline{1}2233\overline{1}1\overline{33} \end{array}$$

This arrangement lives in triple cross surface, decomposed by the curves into 4 digons, 14 tetragons, 3 pentagons, 1 octagon and 1 nonagon. Graphical representations of (tubular neighborhoods of) these arrangements are given in Figure 40.



FIGURE 40. The four martagons on four double pseudolines. Two live in a cross surface and two in a triple cross surface.

Example 14. The 3-chirotope on the 4-set $\{1,2,3,4\}$ with entries $C_{32}(123)$, $C_{32}(124)$, $C_{32}(134)$, $C_{32}(234)$ is not the chirotope of an arrangement because the 3 side cycles indexed by 3 of the entries $C_{32}(123)$, $C_{32}(134)$, $C_{32}(234)$, namely $1\bar{1}22\bar{1}1\overline{22}$, $\bar{4}1\overline{14}4\bar{1}14$, and $\bar{4}2\overline{24}4\bar{2}24$, understood as partial circular orders on the indices $1,2,4$ and their negatives, are incompatibles. Similarly for the 3-chirotope on the 4-set $\{1,2,3,4\}$ with entries $C_{22}(123)$,$C_{22}(423)$, $C_{32}(124), C_{32}(134)$.

Example 15. Example 4 generalizes to any number of indices, i.e., the $C_{64}(ijk)$, where $1 \leq i < j < k \leq n$, is the chirotope of a 3-arrangement on n curves. For $n = 5,6,7,8,9$ we get surfaces of genus $14,21,33,43,58$ decomposed by the curves into 20 2-gons, 1 5-gon, 1 10-gon, 5 16-gons, 1 25-gon for $n = 5$; 30 2-gons, 5 12-gons and 6 20-gons for $n = 6$; 42 2-gons, 1 7-gon, 2 14-gons, 7 24-gons, 1 49-gons for $n = 7$; 56 2-gons, 7 16-gons, 8 28-gons, for $n = 8$; and 72 2-gons, 1 9-gon, 3 18-gons, 3 27-gons, 9 32-gons for $n = 9$.

We come now to the proof that 5-chirotopes are chirotopes of 5-arrangements.

Theorem 27. *Let χ be a 5-chirotope on the indexing set I, let χ^* be its 5-extension, and for $J \in \binom{I}{3,4,5}$ let $D_i(J)$ and $M_i(J)$ be the families of side cycles of the entry $\chi^*(J)$ of χ^*. Then, for any index $i \in I$, there exists*

1. *a unique shuffle D_i of the elementary cycles $jj\overline{jj}$, $j \neq i$, of which the $D_i(J)$, $i \in J \in \binom{I}{3,4,5}$, are subcycles;*
2. *a unique shuffle M_i of the elementary cycles $jj\overline{jj}$, $j \neq i$, of which the $M_i(J)$, $i \in J \in \binom{I}{3,4,5}$, are subcycles.*

Furthermore the two families of cycles D_i and M_i are the families of side cycles of a 5-arrangement of oriented double pseudolines indexed by I with chirotope χ. ☐

Proof. For $X \in \{D, M\}$ and $i \in I$, let $\mathcal{R}_i^X$ be the ternary relation defined on distinct elements $\alpha = i_\alpha j_\alpha \in \{ij, \bar{i}j, i\bar{j}, \overline{ij} \mid j \in I \setminus i\}$, by $(\alpha, \alpha', \alpha'') \in \mathcal{R}_i^X$ if α, α' and α'' appear in this order on the cycle $X_i^*(\{i, j_\alpha, j_{\alpha'}, j_{\alpha''}\})$ and let $\mathcal{B}_i^X$, $i \in I$, be the binary relation defined on distinct elements $\alpha = i_\alpha j_\alpha \in \{ij, \bar{i}j, i\bar{j}, \overline{ij} \mid j \in I \setminus i\}$, by $(\alpha, \alpha') \in \mathcal{B}_i^X$ if α and α' appear in this order in the same prime factor of the prime factor decomposition of $X_i^*(\{i, j_\alpha, j_{\alpha'}\})$. Clearly

1. $\mathcal{R}_i^X$ is well-defined;
2. for every triple $(\alpha, \alpha', \alpha'')$ on has $(\alpha, \alpha', \alpha'') \in \mathcal{R}_i^X$ or (exclusive) $(\alpha, \alpha'', \alpha') \in \mathcal{R}_i^X$;
3. if $(\alpha, \alpha', \alpha'') \in \mathcal{R}_i^X$ then $(\alpha', \alpha'', \alpha) \in \mathcal{R}_i^X$;
4. $\mathcal{R}_i^X$ is transitive, i.e., if $(\alpha, \alpha', \alpha'') \in \mathcal{R}_i^X$ and $(\alpha, \alpha'', \alpha''') \in \mathcal{R}_i^X$ then $(\alpha, \alpha', \alpha''') \in \mathcal{R}_i^X$ (because $X_i^*(i, j_\alpha, j_{\alpha'}, j_{\alpha''})$, $X_i^*(i, j_\alpha, j_{\alpha''}, j_{\alpha'''})$, and $X_i^*(i, j_\alpha, j_{\alpha'}, j_{\alpha''})$ are subcycles of $X_i^*(i, j_\alpha, j_{\alpha'}, j_{\alpha''}, j_{\alpha'''})$);
5. $\mathcal{B}_i^X$ is well-defined;
6. if $(\alpha, \alpha') \in \mathcal{B}_i^X$ and $(\alpha', \alpha'') \in \mathcal{B}_i^X$ then $(\alpha, \alpha'') \in \mathcal{B}_i^X$;
7. if $(\alpha, \alpha'') \in \mathcal{B}_i^X$ and $(\alpha, \alpha', \alpha'') \in \mathcal{R}_i^X$ then (α, α') and $(\alpha', \alpha'') \in \mathcal{B}_i^X$;
8. if $(\alpha, \alpha') \in \mathcal{B}_i^X$ then $(\alpha', \alpha) \in \mathcal{B}_i^{\overline{X}}$.

This proves that the shuffle X_i of the elementary cycles $jj\overline{jj}$, $j \neq i$, given by the ternary relation $\mathcal{R}_i^X$, is the unique shuffle of the elementary cycles $jj\overline{jj}$, $j \neq i$ of which the $X_i(J)$, $i \in J \in \binom{I}{3,4,5}$, are subcycles and that there is a unique block decomposition

$B_{i1}B_{i2}\dots B_{in_i}$, where $B_{im} = \{i_1 j_{m1}\}\{i_2 j_{m2}\}\{i_3 j_{m3}\}\dots\{i_k j_{mk_m}\}$, $1 \leq m \leq n_i$, with $j_{m_l} \neq \pm j_{m_{l'}}$ for all $1 \leq l < l' \leq k_m$, of D_i^* (the one given by the binary relation $\mathcal{B}_i^X$) such that $T_i = B_{i1}^* B_{i2}^* \dots B_{in_i}^*$ where B_{im}^* is the reversal of B_{im}. Since by construction $B_{\otimes j_p}$ is one of the blocks of the block decomposition of $S_{j_{mp}}$ we are done, thanks to Theorem 26. ☐

5.2. **Pumping lemma and mutations.** According to the above discussion we see that the map that assigns to a 5-arrangement its chirotope induces a one-to-one and onto correspondence between the class of isomorphism class of 5-arrangements defined on a given set of indices and the class of 5-chirotopes defined on the same set of indices. We now show that the mutation graph on the space of 5-arrangements defined on a given set of indices is connected. Since a mutation does not change the Euler characteristic of the underlying surface of the 5-arrangement this will prove that 5-arrangements are arrangements of double pseudolines. Actually we prove a stronger result : the space of 4-arrangements defined on a given set of indices is connected.

Theorem 28. *The mutation graph on the space of* 4*-arrangements defined on a given set of indices is connected.* ☐

The proof is based on the following abstractions of the Pumping Lemmas of Section 2.

Lemma 29. *Let τ be a simple* 4*-arrangement of double pseudolines and let $\gamma \in \tau$. Assume that there exists a vertex v of X_τ contained in the crosscap side of γ in the subarrangement of size three composed of γ and of the two double pseudolines crossing at v. Then there exists a triangular* 2*-cell of X_τ with a side supported by γ and a vertex w supported by the crosscap side of γ in the subarrangement composed of γ and of the two double pseudolines crossing at w.* ☐

Lemma 30. *Let τ be a thin* 4*-arrangement of double pseudolines, let $\gamma, \gamma' \in \tau$, $\gamma \neq \gamma'$, and let $M(\gamma,\gamma')$ be one of two two-cells of size* 2 *of the subarrangement $\{\gamma,\gamma'\}$. Assume that there exists a vertex v of X_τ contained in $M(\gamma,\gamma')$ in the subarrangement of size four composed of γ, γ' and of the two double pseudolines crossing at v. Then there exists a triangular* 2*-cell of X_τ contained in $M(\gamma,\gamma')$ with a side supported by γ in the subarrangement composed of γ, γ' and of the two double pseudolines crossing at the vertex w opposite to the side supported by γ.* ☐

Proof of Lemma 29. Without loss of generality we assume that the surfaces S_ν, $\nu \in \binom{\tau}{4}$, intersect pairwise only along their common double pseudolines, i.e., $S_\nu \cap S_{\nu'} = \bigcup \nu''$ where $\nu'' = \nu \cap \nu'$. Let $\mathcal{X}$ be the union of the cross surfaces S_ν as ν ranges over the set of subsets of τ of size at least 2 and at most 4, let X be the cell decomposition of $\mathcal{X}$ induced by τ, let $p : \widetilde{\mathcal{X}} \to \mathcal{X}$ be the two-covering of $\mathcal{X}$ induced by a two-covering of the cross surface, and let $\widetilde{\tau}$ and $\widetilde{X}$ be the "lifts" of τ and X under p, i.e., the connected components of $p^{-1}(\tau)$ and $p^{-1}(X)$. The lifts of γ under p are denoted γ_+ and γ_-. We note that two curves of $\widetilde{\tau}$ have exactly 0 or 2 intersection points depending on whether they are the lifts of the same curve in τ, or not; by convention if B is an intersection point of two crossing curves of $\widetilde{\Gamma}$ then the second one is denoted B_* (see Figure 41).

Define a γ*-curve* as a maximal subcurve of some $\widetilde{\gamma}' \in \widetilde{\Gamma}$, $\gamma' \neq \gamma$, contained in the cylinder $\widetilde{M}$, where M is the Möbius strip bounded by γ in $S_{\{\gamma,\gamma'\}}$. We observe that



FIGURE 41. (Top) An arrangement of two double pseudolines γ, γ' in $S_{\{\gamma,\gamma'\}}$ (represented here by the polygonal schema $\alpha\alpha = 1$) and its lift in $\widetilde{S}_{\{\gamma,\gamma'\}}$ (represented here by the polygonal schema $\alpha\alpha^{-1} = 1$). (Bottom) The cell decomposition $C(B, B')$ of the cylinder strip$(\{B, B'\})$ induced by the γ-curves curve$_\gamma(B)$ and curve$_\gamma(B')$ depends only on their number of intersection points.

there are two γ-curves per $\widetilde{\gamma}' \in \widetilde{\Gamma}$, $\gamma' \neq \gamma$, and a γ-curve has an endpoint on γ_+ and the other one on γ_-; we denote by curve$_\gamma(B)$ the γ-curve with endpoint B on γ_+.

Given a set $\mathcal{B}$ of at most 4 vertices on the curve γ^+ we denote by strip$(\mathcal{B})$ the cylinder $\widetilde{M_J}$ where J is the set of curves in τ supporting the γ-curves curve$_\gamma(B)$, $B \in \mathcal{B}$, augmented with γ and where M_J is the Möbius strip bounded by γ in S_J. We say that strip$(\mathcal{B})$ is the cylinder defined by $\mathcal{B}$ or by curve$_\gamma(\mathcal{B})$.

Two γ-curves curve$_\gamma(B)$ and curve$_\gamma(B')$ induce a cell decomposition $C(B, B')$ of the cylinder strip$(\{B, B'\})$ which depends only on their number (0, 1 or 2) of intersection points; this cell decomposition is composed of two quadrilaterals or is composed of two triangles and a hexagon or is composed of a digon, two triangles, and an octagon — according to the number 0, 1 or 2 of intersection points—put together as depicted in Figure 41.

Define a γ-*triangle* as a triangular face of the cell decomposition of the cylinder C defined by two crossing γ-curves with a side supported by γ_+; the vertex of a γ-triangle not on γ_+ is called its *apex* and the side of a γ-triangle supported by γ_+ called its *base side.* A γ-triangle is said to be *admissible* if one of its two sides with endpoint its apex is an edge of $\widetilde{X}$. The interior of the base side of a γ-triangle T is denoted $\mathrm{bs}_\gamma(T)$ and its complement in γ_+ is denoted $\overline{\mathrm{bs}}_\gamma(T)$.

Let $T = ABB'$ be an admissible γ-triangle with apex A and edge side AB, and let A' be the vertex of $\widetilde{X}$ that follows B' on the side $B'A$ of T. Then A' is the apex of an admissible γ-triangle $T' = A'B'B''$ with edge side $A'B'$.

A simple use of the Jordan curve theorem leads to the three following lemmas that control the relative positions of the base sides of $T, T', T'', T''', \ldots$. (See Figures 42 and **??** for an illustration).



$B'' \in \overline{\mathrm{bs}}_\gamma(T)$ $B'' \in \mathrm{bs}_\gamma(T)$

$\mathrm{strip}(\{B, B'\})$ $\mathrm{strip}(\{B, B', B''\})$ $\mathrm{strip}(\{B, B', B''\})$

FIGURE 42.

Lemma 31. *Assume that* $T = T'$. *Then* T *is a triangular two-cell of* $\widetilde{X}$. ☐

Lemma 32. *Assume that* $T \neq T'$, *that* $B'' \in \overline{\mathrm{bs}}_\gamma(T)$. *Then*

(1) $\mathrm{curve}_\gamma(B')$ *and* $\mathrm{curve}_\gamma(B'')$ *cross twice (at* A' *and* A'_**) on the side* $B'A$ *of* T,
(2) $\mathrm{bs}_\gamma(T)$ *and* $\mathrm{bs}_\gamma(T')$ *are interior disjoint,*
(3) B'_* *and* $B''_* \in \overline{\mathrm{bs}}_\gamma(T) \cap \overline{\mathrm{bs}}_\gamma(T')$, *and*
(4) walking along $\overline{\mathrm{bs}}_\gamma(T) \cap \overline{\mathrm{bs}}_\gamma(T')$ *from* B'' *to* B *we encounter successively the points* B''_* *and* B'_*. ☐

Lemma 33. *Assume that* $T \neq T'$, *that* $B'' \in \mathrm{bs}_\gamma(T)$. *Then (1)* $\mathrm{curve}_\gamma(B'')$ *crosses the side* $B'A$ *of* T *exactly once (at* A'*) and (2)* $\mathrm{bs}_\gamma(T')$ *is contained in* $\mathrm{bs}_\gamma(T)$. ☐

An admissible γ-triangle $\Delta = XYY'$ with apex X and edge side XY is said to enclose T if the three following conditions are satisfied

(1) the base side of T is included in the base side of Δ and walking along the base side of Δ from Y to Y' we encounter B' before B, as illustrated in the left part of Figure 43 (we allow $B = Y'$);
(2) the arrangement of the three γ-curves $\mathrm{curve}_\gamma(Y'), \mathrm{curve}_\gamma(B), \mathrm{curve}_\gamma(B')$ is, up to homeomorphism, the one depicted in the middle part of Figure 43;



strip($\{Y, Y'\}$) strip($\{B, B', Y'\}$) strip($\{B, B', Y\}$)

Figure 43.

(3) the arrangement of the three γ-curves $\mathrm{curve}_\gamma(Y), \mathrm{curve}_\gamma(B), \mathrm{curve}_\gamma(B')$ is, up to homeomorphism, the one depicted in the right part of Figure 43.

Lemma 34. *Assume that* $T \neq T'$, *that* $B'' \in \overline{\mathrm{bs}}_\gamma(T)$ *and that there is an admissible* γ*-triangle* Δ *that encloses* T. *Then* Δ *encloses* T'. ☐



$B'' \in \overline{\mathrm{bs}}_\gamma(T)$

strip($\{B, B', B''\}$)

Figure 44.

Consider now the sequence of admissible γ-triangles $T_0, T_1, T_2, \ldots$ defined inductively by $T_0 = T$ and $T_{k+1} = T'_k$ for $k \geq 0$. A simple combination of Lemmas 32, 33, and 34 leads to the conclusion that the sequence T_k is stationary. According to Lemma 31 the lemma follows. ☐

Proof of Lemma 30. The proof using similar ideas to the proof of the previous lemma with a much simpler case analysis we omit it. ☐

Proof of Theorem 28. Clearly any 4-arrangement is connected, via a finite sequence of splitting mutations, to a simple 4-arrangement. By a repeated application of Lemma 29 we see that any simple 4-arrangement is connected, via a finite sequence of mutations, to a simple thin 4-arrangement. Then by a repeated application of Lemma 30 we see that any simple thin 4-arrangement is connected, via a finite sequence of mutations, to the double version of cyclic arrangement of pseudolines. ☐

Remark 2. Our axiomatic characterization of the class of chirotopes of double pseudoline arrangements asserts that the full list of indexed arrangements on at most five oriented double pseudolines (on a given set of five indices) can be viewed as an axiom systems for the class of double pseudoline arrangements. Since the number of arrangements on five double pseudolines is in the order of several hundred millions [19], this list seems hardly computationally workable for the generation of arrangements with prescribed properties. However, since the class of 4-arrangements coincides with the class of 5-arrangements, it is possible to replace the huge list of axioms involving exactly five double pseudolines by a single axiom that simply states that certain ternary relations—the $\mathcal{R}_i^X$ of Section 5.1—are total circular orders. This drastic reduction in the number of axioms opens the door to the generation of double pseudoline arrangements with prescribed properties using, for example, satisfiability solvers as proposed in [42, 41]. In particular it will be interesting to generate the arrangements that maximize the number of connected components of the intersection of the crosscap sides of the double pseudolines. If we think of the arrangement as the dual of a configuration of pairwise disjoint convex bodies, this means that we seek the configurations with the maximal number of connected components of transversals.

Remark 3. Other proofs of our axiomatization characterization of double pseudoline arrangements are conceivable. One can try to proceed by induction as it is done in the recent alternative proof of the Folkman and Lawrence axiomatic characterization of arrangements of pseudohyperplanes given in [4, 5, 3] or, back to the late seventies, by establishing suitable shelling properties of the face poset of 4-arrangements as done by Folkman and Lawrence for their axiomatic characterization of arrangements of pseudohyperplanes [20, 2].

6. An extension and a refinement

In this sixth and next to last section we discuss *arrangements of pseudocircles*, *crosscap or Möbius arrangements* and the *fibrations* of the latter. The material on fibrations was partially motivated by the question raised by J.-E. Goodman and R. Pollack in [26, Remark 20] about the realizability of their so-called double-permutation sequences by families of pairwise disjoint convex bodies of affine planes.

Define an *oval* as the boundary of a convex body of a projective plane and the *dual of an oval* as the set of lines touching the oval but not its disk side.

An *arrangement of pseudocircles* is a finite family of pseudocircles embedded in a cross surface, with the property that its subfamilies of size two are homeomorphic to the dual arrangement of two points, two disjoint ovals, or one point and one oval which are not incident. Observe that arrangements of pseudocircles extend both arrangements of pseudolines and arrangements of double pseudolines. Figure 45 depicts representatives of the isomorphism classes of arrangements of two pseudocircles. The *order* of an arrangement of pseudocircles is defined as the isomorphism class of the poset of the bicolored curves of the arrangement ordered by reverse inclusion of their disk sides where by convention a pseudoline is colored in blue ($\bullet$) and a double pseudoline is colored in red ($\bigcirc$). For example the orders of the arrangements depicted in the above figure are $\{\bigcirc \rightarrow \bigcirc\}, \{\bigcirc, \bigcirc\}, \{\bullet \rightarrow \bigcirc)\}, \{\bullet, \bullet\}$, and $\{\bullet, \bigcirc\}$. According to Theorem 1 examples of arrangements of pseudocircles are given by the dual arrangements of finite families of pairwise disjoint ovals and points of projective planes. Minor adaptations in our proof of the Pumping Lemma for arrangements of double pseudolines lead to the following Pumping Lemma for arrangements of pseudocircles. We denote by $\mathcal{D}(\gamma)$ the disk side of a double pseudoline γ.



Figure 45. Arrangements on two pseudocircles.

Lemma 35 (Pumping Lemma for Arrangements of Pseudocircles)**.** *Let Γ be a simple arrangement of pseudocircles, let $\gamma \in \Gamma$ be a double pseudoline of Γ, let Γ' be the set*

$\gamma' \in \Gamma$ such that $\mathcal{D}(\gamma') \supset \mathcal{D}(\gamma)$, and let M be the trace on the crosscap side of γ of the 2-cell of the arrangement Γ' that contains $\mathcal{D}(\gamma)$. Assume that there is a vertex of the arrangement Γ lying in M. Then there is a triangular two-cell of the arrangement Γ contained in M with a side supported by γ. □

Then minor variations in our proofs of Theorems 2, 3, and 4 based on the above Pumping Lemma lead to direct extensions of these theorems modulo the following elementary dictionnary:

arrangements of double pseudolines ⟷ arrangements of pseudocircles
convex bodies ⟷ ovals and points
size ⟷ order

In particular

Theorem 36. *Any arrangement of pseudocircles is isomorphic to the dual family of a finite family of pairwise disjoint ovals and points of a projective plane.* □

For the numbers of isomorphism classes of simple arrangements with trivial order ({●, . . . , ●, ○, . . . , ○}) on at most five curves we refer to [19], where these arrangements are called mixed arrangements.

We now discuss a refinement of Theorem 36 in the context of *crosscap or Möbius arrangements* and their *fibrations.* Let $\mathcal{M}$ be a Möbius strip, let $\overline{\mathcal{M}} = \mathcal{M} \cup \{\infty\}$ be its one-point compactification, and recall that $\overline{\mathcal{M}}$ is a cross surface. Define a *arrangement of pseudocircles in* $\mathcal{M}$ as a arrangement of pseudocircles in $\overline{\mathcal{M}}$ with the property that the intersection of the disk sides of its pseudolines and double pseudolines is nonempty and contains the point at infinity; define a *fibration* of an arrangement of pseudocircles Γ in $\mathcal{M}$ as a sup-arrangement Γ' of Γ in $\overline{\mathcal{M}}$ composed of the pseudocircles of Γ and of the pseudolines of a pencil of pseudolines through the point at infinity with the property that any pseudoline of the pencil goes through a vertex of Γ and any vertex of Γ is incident to a pseudoline of the pencil. Figure 46 shows an arrangement of two double pseudolines in a Möbius strip and representatives of its three possible isomorphism classes of fibrations. According to Theorem 36 we see easily that any arrangement of pseudocircles Γ in $\mathcal{M}$ is the dual arrangement of a family of pairwise disjoint ovals and points of a projective plane $\mathcal{G}$ with line space $\overline{\mathcal{M}}$, with the property that the line at infinity ∞ avoids the ovals



Figure 46. An arrangement of two double pseudolines in a Möbius strip with representatives of its three isomorphism classes of fibrations.

and the points of the family. Consequently the family Γ' composed of the pseudocircles of Γ and of the dual pseudolines of the intersection points of the line at infinity with the vertices of Γ is a fibration of Γ. This fibration is denoted $\Gamma^{\mathcal{G}}$. Conversely let Γ' be a fibration of Γ. Does there exist a projective plane $\mathcal{G}$ such that $\Gamma' = \Gamma^{\mathcal{G}}$? Applying Theorem 36 to Γ' we obtain a positive answer to that question.

Theorem 37. *Let $\mathcal{M}$ be a Möbius strip and let Γ' be a fibration of an arrangement of pseudocircles Γ in $\mathcal{M}$. Then Γ is the dual family of finite family of pairwise disjoint ovals and points of a projective plane $\mathcal{G}$ with line space $\overline{\mathcal{M}}$ such that $\Gamma' = \Gamma^{\mathcal{G}}$.* □

In particular the above theorem answers positively the question raised by J.-E. Goodman and R. Pollack in [26, Remark 20] about the realizability of their so-called double-permutation sequences by families of pairwise disjoint convex bodies of real two-dimensional affine geometries. Indeed double-permutation sequences are simply coding of isomorphism classes of our fibrations. For example the double permutation sequences coding the three isomorphism classes of fibrations of an arrangement of two double pseudolines indexed by $\{1, 2\}$ are the following

$$(*) \qquad \begin{bmatrix} 2 & 2 & 2 & 1 & 1 \\ 2 & 1 & 1 & 2 & 1 \\ 1 & 2 & 1 & 1 & 2 \\ 1 & 1 & 2 & 2 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 2 & 1 & 1 \\ 2 & 1 & 2 & 1 \\ 1 & 2 & 1 & 2 \\ 1 & 1 & 2 & 2 \end{bmatrix}, \begin{bmatrix} 2 & 2 & 1 & 1 & 1 \\ 2 & 1 & 2 & 2 & 1 \\ 1 & 2 & 2 & 1 & 2 \\ 1 & 1 & 1 & 2 & 2 \end{bmatrix}.$$

We refer to [26] for the precise definition of double permutation sequences.

We conclude by the statements of the counterparts of Theorems 2 and 4 in the context of arrangements in Möbius strips.

Theorem 38. *Let $\mathcal{M}$ be a Möbius strip. Any two arrangements of pseudocircles in $\mathcal{M}$ of the same order are homotopic in $\mathcal{M}$ via a finite sequence of mutations followed by an isotopy.* □

An arrangement of oriented pseudocircles is termed *acyclic* if the orientations of the pseudocircles are coherent, in the sense that the pseudocircles are oriented according to the choice of a generator of the (infinite cyclic) fundamental group of the underlying Möbius strip.

Theorem 39. *The map that assigns to the isomorphism class of an indexed Möbius arrangement of oriented pseudocircles its chirotope is one-to-one and its range is the set of maps χ defined on the set of triples of a finite set I such that for every 3-, 4-, and 5-subset J of I the restriction of χ to the set of triples of J is the chirotope of an indexed Möbius arrangement of oriented pseudocircles. Furthermore the same result holds for the class of acyclic Möbius arrangements of pseudocircles.* □

Note that our homotopy theorem for Möbius arrangements provides an algorithm to enumerate the isomorphism classes of Möbius arrangements by a traversal of the associated mutation graphs. We have implemented this algorithm to enumerate the isomorphism classes of the simple Möbius arrangements of double pseudolines. Preliminary counting results (confirmed by two independent implementations) are reported in the following table

n	1	2	3	4
a_n	1	1	118	541820
b_n	1	1	22	22620
c_n	1	1	16	11502
d_n	1	1	12	5955

– here the index n refers to the number of double pseudolines, a_n is the number of isotopy classes of simple indexed arrangements of double pseudolines, b_n is the number of isotopy classes of simple arrangements of double pseudolines, c_n is the number of isomorphism classes simple arrangements of double pseudolines, and d_n is the number of isomorphism classes of simple arrangements of double pseudolines considered as projective arrangements. Representatives of the sixteen isomorphism classes of simple Möbius arrangements of three double pseudolines are depicted in Figure 47. There are 531 chirotopes on a given indexing set of size 3 and among these 531 chirotopes 118 are simple chirotopes [19]. Representatives of these 118 chirotopes are implicitly depicted in Figure 47 : each diagram represents an isomorphism class of non indexed arrangement of three double pseudolines. Each diagram is labeled at its left bottom corner with the



Figure 47.

2-sequence of its numbers of 2-cells of size 2 and 3 and is labeled at its right bottom corner with the size of its automorphism group.

It is interesting to mention that the canonical embedding of Lemma 6 can be extended, in the case of Möbius arrangements, to the whole class of simple and nonsimple arrangements as explain below.

Define a *pencil* of double pseudolines as a Möbius arrangement of double pseudolines with the property that any of its subarrangements has only two *external* vertices, i.e., only two vertices in the boundary of the two-cell that contains the point at infinity of the one-point compactification of the underlying Möbius strip. Figure 48 shows pencils of two, three, four and five double pseudolines; it is not hard to see that the isotopy class of a pencil of double pseudolines depends only on its number of double pseudolines. Now a Möbius arrangement of double pseudolines is termed *thin* if any of its subarrangements whose double pseudolines have associated Möbius strips with nonempty intersection is a pencil of double pseudolines—to put it differently: a Möbius arrangement of double pseudolines is thin if the crosscap sides of its double pseudolines are free of external vertices—and a Möbius arrangement of double pseudolines Γ^* is termed a *double* of a Möbius pseudoline arrangement Γ if there exists a one-to-one correspondence between Γ and Γ^* such that



Figure 48. Pencils of 2, 3, 4, and 5 double pseudolines.

1. any pseudoline of Γ is contained in the crosscap side of its corresponding double pseudoline in Γ^*, and
2. any subarrangement of Γ^* is a pencil of double pseudolines if and only if the corresponding subarrangement of Γ is a pencil of pseudolines.

In this Möbius setting Lemma 6 can be read as follow.

Lemma 40. *Let $\mathcal{M}$ be a Möbius strip. The map that assigns to an arrangement of pseudolines in $\mathcal{M}$ its set of double versions induces a one-to-one and onto correspondence between the set of isotopy classes of pseudoline arrangements in $\mathcal{M}$ and the set of isotopy classes of thin double pseudoline arrangements in $\mathcal{M}$.* ☐

7. Conclusion and open problems

We have introduce the notion of arrangements of double pseudolines as a combinatorial abstraction of families of pairwise disjoint convex bodies of projective planes and we have extended to that setting well-known fundamental properties of arrangements of pseudolines.

Several open questions are raised by our work. We mention below three of them.

The cell poset of a simple arrangement of pseudolines presented by its chirotope is computable in optimal quadratic time and linear working space (cf. [17]). Can we achieve similar bounds for arrangements of double pseudolines presented by their chirotopes? Progress in this direction using the notion of pseudotriangulation are reported in our companion paper [28].

There is a closed formula, due to R. Stanley, counting the number r_n of wiring representations of simple Möbius arrangements of n pseudolines, namely

$$r_n = \frac{\binom{n}{2}!}{(2n-3)(2n-5)^2(2n-7)^3 \cdots 5^{n-3}3^{n-2}}.$$

Most of the proofs of this formula if not all are based on connections between standard Young Tableaux, reduced words and arrangements of pseudolines (see [43, 16, 34, 29, 18]). Are there similar connections for Möbius arrangements of double pseudolines? Is there any similar formula counting the number of wiring representations of simple Möbius arrangements of double pseudolines?

Say that an arrangement of n double pseudolines is realizable if it is the dual of a family of n disjoint discs of the standard two-dimensional projective plane $\mathcal{P}_2(\mathbb{R})$. In that case one can think of the arrangement as the trace on the unit sphere of $\mathbb{R}^3$ of a centrally symmetric affine arrangement of $2n$ planes with the property that the distance to the origin of any line defined as the intersection of d of these planes is less than 1. The open question is then the following : Is any arrangement of double pseudolines the trace on the unit sphere of a centrally symmetric affine arrangement of pseudoplanes in $\mathbb{R}^3$?

Arrangements of double pseudolines are dual families of families of pairwise disjoint convex bodies of projective planes. What is the smallest example which is not realizable as the dual of a family of pairwise disjoint disks of the standard projective plane $\mathcal{P}_2(\mathbb{R})$? (By a disk we mean a convex body whose boundary is a circle, i.e., the intersection of the unit sphere of $\mathbb{R}^3$ with an affine plane.)

Arrangements of double pseudolines generalize arrangements of pseudolines. What are the similar generalizations for arrangements of pseudohyperplanes of dimensions 4, 5, etc.? A generalization that comes naturally to mind defines (1) a double pseudohyperplane as the image of the hypersurface $x_1 = \pm 1/\sqrt{2}$ of the projective space $\mathbb{RP}^d$ (defined as the quotient of the unit sphere of $\mathbb{R}^{d+1}$ under the antipodal map) under a self-homeomorphism of $\mathbb{RP}^d$, and (2) an arrangement of double pseudohyperplanes as a finite family of double pseudohyperplanes with the property that its subfamilies of size d are the images of the arrangement composed of the d hypersurfaces $x_i = \pm 1/\sqrt{2d}$, $i = 1, 2, \ldots, d$, under a self-homeomorphism of $\mathbb{RP}^d$; two questions show up naturally : (1) Does the isomorphism class of a (indexed and oriented) double pseudohyperplane arrangement depend only on its chirotope, i.e., the family of isomorphism classes of its subarrangements of size $d+1$? (2) Does the class of chirotopes of double pseudohyperplane arrangements coincide with the class of maps that assigns to each $d+1$-subset of indices an isomorphism class of arrangements of double pseudohyperplanes indexed by that $d+1$-subset and whose restrictions to the sets of $d+1$-subsets of $d+3$-subsets of indices are chirotopes of double pseudohyperplane arrangements?

Acknowledgments.

We thank Günter Rote for fruitful discussions, concerning the research reported in this paper, during his visit at the Ecole Normale Supérieure in October 2005, we thank the Mezzenile research group for its constant support and for providing us an independent implementation of the enumeration algorithm described in this paper, and we thank Ricky Pollack for his constructive criticism of a first draft of that paper.

References

[1] M. Berger. *Géométrie: Convexes et Polytopes, Polyèdres Réguliers, Aires et Volumes*, volume 3. Cedic/Fernand Nathan, 1978.

[2] A. Björner, M. Las Vergnas, B. Sturmfels, N. White, and G. M. Ziegler. *Oriented Matroids*. Cambridge University Press, Cambridge, 2 edition, 1999.

[3] Jürgen Bokowski. *Computational Oriented Matroids*. Cambridge, 2006.

[4] Jürgen Bokowski, Simon King, Suzanne Mock, and Ileana Streinu. The topological representation of oriented matroids. *Disc. Comput. Geom.*, 33(4):645–668, 2005. `http://www.springerlink.com/content/q2034875503tt61u/fulltext.pdf`.

[5] Jürgen Bokowski, Suzannne Mock, and Ileana Streinu. On the Folkman-Lawrence topological representation theorem for oriented matroids of rank 3. *Europ. J. Comb.*, 22(5):601–615, 2001.

[6] H. Busemann and H. Salzmann. Metric collineations and inverse problems. *Math. Zeit.*, 87(1):214–240, 1965.

[7] J. Cantwell. Geometric convexity. I. *Bull. Inst. Math. Acad. Sinica*, 2:289–307, 1974.

[8] J. Cantwell. Geometric convexity. II: Topology. *Bull. Inst. Math. Acad. Sinica*, 6:303–311, 1978.

[9] J. Cantwell and D. C. Kay. Geometric convexity. III: Embeddings. *Trans. Amer. Math. Soc.*, 246:211–230, 1978.

[10] R. Cordovil. Sur les matroides orientés de rang 3 et les arrangements de pseudodroites dans le plan projectif réel. *European J. Combinatorics*, 3:307–318, 1982.

[11] J. de Groot and H. de Vries. Convex sets in projective geometry. *Compositio Mathematica*, 13:113–118, 1958.

[12] David Dekker. Convex regions in projective n-space. *Amer. Math. Monthly*, 62(6):430–431, 1955.

[13] R. Dhandapani, J. E. Goodman, A. Holmsen, R. Pollack, and S. Smorodinsky. Convexity in topological affine planes. *Disc. Comput. Geom.*, 38:243–257, 2007.

[14] J-P. Doignon. Caractérisations d'espaces de Pasch-Peano. *Bull. Acad. Roy. Belg. Cl. Sci.*, 62(5):679–699, 1976.

[15] Milton Drandell. Generalized convex sets in the plane. *Duke Math. J.*, 19:537–547, 1952.

[16] P. Edelman and C. Greene. Balanced tableaux. *Adv. Math.*, 63:42–99, 1987.

[17] H. Edelsbrunner and Leonidas J. Guibas. Topologically sweeping an arrangement. *J. Comput. Syst. Sci.*, 38(1):165–194, 1989. Corrigendum in 42 (1991), 249–251.

[18] Stefen Felsner. The skeleton of a reduced word and a correspondence of Edelman and Greene. *The Electronic Journal of Combinatorics*, 8(1):R8, 2001.

[19] Julien Ferté, Vincent Pilaud, and Michel Pocchiola. On the number of simple arrangements of five double pseudolines. *Disc. Comput. Geom.*, 45(2):279–302, 2011. Special issue on Transversal and Helly-type Theorems in Geometry, Combinatorics and Topology. A preliminary version appeared in the abstracts 18th fall Workshop Comput. Geom., Troy, 2008 (FWCG 2008). DOI: 10.1007/s00454-010-9298-4.

[20] J. Folkman and J. Lawrence. Oriented matroids. *J. Combinatorial Theory Ser. B*, 25:199–236, 1978.

[21] Georges K. Francis and Jeffrey R. Weeks. Conways's ZIP proof. *Amer. Math. Monthly*, 106(5):393–399, May 1999.

[22] J. E. Goodman. Proof of a conjecture of Burr, Grünbaum and Sloane. *Discrete Math.*, 32:27–35, 1980.

[23] J. E. Goodman and R. Pollack. Semispaces of configurations, cell complexes of arrangements. *J. Combin. Theory Ser. A*, 37:257–293, 1984.

[24] J. E. Goodman, R. Pollack, R. Wenger, and T. Zamfirescu. Arrangements and topological planes. *Amer. Math. Monthly*, 101(9):866–878, November 1994.

[25] Jacob E. Goodman. Pseudoline arrangements. In Jacob E. Goodman and Joseph O'Rourke, editors, *Handbook of Discrete and Computational Geometry*, chapter 5, pages 97–128. Chapman & Hall/CRC, 2004.

[26] J.E. Goodman and R. Pollack. The combinatorial encoding of disjoint convex sets in the plane, and a generalization of the edelsbrunner-sharir transversal theorem, 2004.

[27] B. Grünbaum. *Arrangements and Spreads*. Regional Conf. Ser. Math. American Mathematical Society, Providence, RI, 1972.

[28] Luc Habert and Michel Pocchiola. Computing pseudotriangulations via branched covering. Technical Report arXiv:1102.0151v1 [cs.CG] 1 Fev. 2011, arXiv, 2011.

[29] M. Haiman. Dual equivalence with applications, including a conjecture of proctor. *Discrete Math.*, 99:79–113, 1992.

[30] Allan Hatcher. *Algebraic Topology.* Cambridge University Press, 2001.

[31] W. Holsztynski, J. Mycielski, G. C. O'Brien, and S. Swierczkowski. Closed curves on spheres. *Amer. Math. Monthly*, 115(2):149–153, 2008.

[32] Robert C. James. Combinatorial topology of surfaces. In Ann K. Stehney, Tilla K. Milnor, Joseph E. D'Atri, and Thomas F. Banchoff, editors, *Selected papers on geometry*, volume 4 of *The Raymond W. Brink Selected Mathematical Papers*, pages 79–114. The Mathematical Association of America, 1979. Reprint from Mathematics Magazine, vol. 20 (1955), pp. 1–39.

[33] David C. Kay and Eugene W. Womble. Axiomatic convexity theory and relationships between the Carathéodory, Helly, and Radon numbers. *Pacific J. Math.*, 38(2):471–485, 1971.

[34] Alain Lascoux and Marcel-Paul Schützenberger. Structure de Hopf de l'anneau de cohomologie et de l'anneau de grothendiek d'une variété de drapeaux. *Comptes Rendus des séances de l'Académie des Sciences, Série I*, 295:629–633, 1982.

[35] William S. Massey. *A Basic Course in Algebraic Topology.* Springer-Verlag, 1991.

[36] Burkard Polster and Günter Steinke. *Geometries on Surfaces.* Number 84 in Encyclopedia of Mathematics and its applications. Cambridge, 2001.

[37] G. Ringel. Teilungen der Ebene durch Geraden oder topologische Geraden. *Math. Zeitschrift*, 64:79–102, 1956.

[38] G. Ringel. Über Geraden in allgemeiner Lage. *Elemente der Math.*, 12:75–82, 1957.

[39] H. R. Salzmann. Topological planes. *Adv. Math.*, 2:1–60, 1968.

[40] Helmut Salzmann, Dieter Betten, Theo Grundhöfer, Hermann Hähl, Rainer Löwen, and Markus Stroppel. *Compact projective planes.* Number 21 in De Gruyter expositions in mathematics. Walter de Gruyter, 1995.

[41] Lars Schewe. Generation of oriented matroids using satisfiability solvers. In *ICMS*, volume 4151 of *LNCS*, pages 216–218, 2006. `http://www.springerlink.com.gate6.inist.fr/content/7228340602700123/fulltext.pdf`.

[42] Lars Schewe. Non-realizable minimal vertex triangulations of surfaces: showing non-realizability using oriented matroids and satisfiability solvers. Technical Report arXiv:0801.2582v1 [math.MG] 16 Jan 2008, arXiv, January 2008.

[43] R. Stanley. On the number of reduced decompositions of elements of Coxeter groups. *European J. Combin.*, 5:359–372, 1984.

[44] Bernd Sturmfels and Günter M. Ziegler. Extension spaces of oriented matroids. *Disc. Comput. Geom.*, 10:23–45, 1993.

Appendix A. Arrangements of pseudolines

An *arrangement of pseudolines* is a finite set of pseudolines living in the same cross surface with the property that any two pseudolines intersect in exactly one point and the *isomorphism class* of an arrangement of pseudolines is its set of homeomorphic images. Figure 49 depicts representatives of the (full list of) isomorphism classes of arrangements of at most five pseudolines. Each arrangement is labeled at its right bottom corner with the size of its automorphism group.



Figure 49. Representatives of the isomorphism classes of arrangements of one, two, three, four and five pseudolines.

The question of understanding the isomorphism relation between arrangements of pseudolines was addressed and solved by Folkman and Lawrence [20]—in the broader context of arrangements of pseudohyperplanes—essentially as indicated in the following theorem where the term *chirotope* applied to an indexed arrangement of oriented pseudolines means the map that assigns to each triple of indices of the arrangement the isomorphism class of the subarrangement indexed by this triple. (The case of simple arrangements was already studied and solved in a somewhat different but equivalent formulation by Ringel [37, 38].)

Theorem 41 ([20])**.** *The map that assigns to the isomorphism class of an indexed arrangement of oriented pseudolines its chirotope is one-to-one and its range is the set* *of map χ defined on the set of triples of a finite set I such that for every 3-, 4-, and 5-subset J of I the restriction of χ to the set of triples of J is the chirotope of an arrangement of oriented pseudolines.* □

The above theorem can be completed by a comprehensive description of the indexed and oriented arrangements on 3, 4 and 5 pseudolines as we now explain. Let X be an arrangement of pseudolines, let X_* be an indexed and oriented version of X, say indexed by the indexing set I, and extend X_* to the set of negatives of the elements of I by assigning to the negative of $i \in I$ the reoriented version of the pseudoline assigned to i. Let G_I be the group of permutations of the signed indices which are compatible with the operation of taking the negative and let G_X be the group of automorphisms of X. Clearly the map that assigns to $\sigma \in G_X$ its conjugate $X_*^{-1}\sigma X_* \in G_I$ under X_* is a monomorphism of G_X into G_I. Thus we can see G_X as a subgroup G_{X_*} of G_I and the number of distinct indexed and oriented versions of X is the index $[G_I : G_{X^*}]$ of G_{X_*} in G_I. In the sequel we use the notation $X(\sigma)$ for the arrangement $X_*\sigma$, $\sigma \in G_I$; hence $X(1) = X_*$, where 1 is the unit of G_I.

Example 16. Figure 50 depicts an arrangement H on 5 pseudolines, its first barycentric subdivision, and one of its indexed and oriented version H_* on the indexing set $\{1,2,3,4,5\}$. The group G_H is a dihedral group of order 8 generated, for example, by the automorphism σ_{12} that exchanges the flags numbered 1 and 2 in the figure and the automorphism σ_{18} that exchanges the flags numbered 1 and 8 in the figure. Thus the number of distinct indexed and oriented versions of H is $5!2^5/8 = 480$. The group G_{H_*} is generated by the permutations $1\overline{54}23$ and $\overline{1}32\overline{45}$ which correspond to the automorphisms σ_{12} and σ_{18}, respectively.



Figure 50.

Figure 51 depicts one indexed and oriented version of each of the isomorphism classes of arrangements of three and four pseudolines; each diagram is labeled at its right bottom corner by its number of distinct reindexations (on a given set of indices) and reorientations. Thus the arrangement A has 2 distinct indexed and oriented versions. The group G_{A_*} is S_4 generated, for example, by the permutations $\overline{1}32$, $\overline{12}3$ and $2\overline{31}$ and its 2 cosets



$A(123)$ 2 $B(123)$ 4

$C(1234)$ 16 $D(1234)$ 32 $E(1234)$ 24

Figure 51. Isomorphism classes of arrangements of three and four oriented pseudolines indexed by $1, 2, 3, 4$.

are

$$G_{A_*} = \left\{ \begin{array}{ccc} 123 & 231 & 312 \\ \overline{12}3 & \overline{23}1 & \overline{31}2 \\ \overline{1}2\overline{3} & \overline{2}3\overline{1} & \overline{3}1\overline{2} \\ 1\overline{23} & 2\overline{31} & 3\overline{12} \\ 21\overline{3} & 32\overline{1} & 13\overline{2} \\ 3\overline{2}1 & 1\overline{3}2 & 2\overline{1}3 \\ \overline{1}32 & \overline{2}13 & \overline{3}21 \\ \overline{213} & \overline{321} & \overline{132} \end{array} \right\}, \qquad (213)G_{A_*} = \left\{ \begin{array}{ccc} 213 & 321 & 132 \\ \overline{21}3 & \overline{32}1 & \overline{13}2 \\ \overline{2}1\overline{3} & \overline{3}2\overline{1} & \overline{1}3\overline{2} \\ 2\overline{13} & 3\overline{21} & 1\overline{32} \\ 12\overline{3} & 23\overline{1} & 31\overline{2} \\ 3\overline{1}2 & 1\overline{2}3 & 2\overline{3}1 \\ \overline{2}31 & \overline{3}12 & \overline{1}23 \\ \overline{123} & \overline{231} & \overline{312} \end{array} \right\}.$$

Similarly the number of distinct indexed and oriented versions of the arrangement B is 4. The group G_{B_*} is $S_3 \times \mathbb{Z}_2$, generated for example by the permutations $231, 213, \overline{123}$ and its 4 cosets are

$$G_{B_*} = \left\{ \begin{array}{ccc} 123 & 231 & 312 \\ 213 & 321 & 132 \\ \overline{123} & \overline{231} & \overline{312} \\ \overline{213} & \overline{321} & \overline{132} \end{array} \right\}, (\overline{1}23)G_{B_*} = \left\{ \begin{array}{ccc} \overline{1}23 & 23\overline{1} & 3\overline{1}2 \\ 2\overline{1}3 & 32\overline{1} & \overline{1}32 \\ 1\overline{23} & \overline{23}1 & \overline{3}1\overline{2} \\ \overline{2}1\overline{3} & \overline{32}1 & 1\overline{32} \end{array} \right\},$$

$$(1\overline{2}3)G_{B_*} = \left\{ \begin{array}{ccc} 1\overline{2}3 & \overline{2}31 & 31\overline{2} \\ \overline{2}13 & 3\overline{2}1 & 13\overline{2} \\ \overline{1}2\overline{3} & 2\overline{31} & \overline{31}2 \\ 2\overline{13} & \overline{3}2\overline{1} & \overline{13}2 \end{array} \right\}, (12\overline{3})G_{B_*} = \left\{ \begin{array}{ccc} 12\overline{3} & 2\overline{3}1 & \overline{3}12 \\ 21\overline{3} & \overline{3}21 & 1\overline{3}2 \\ \overline{12}3 & \overline{2}3\overline{1} & 3\overline{12} \\ \overline{21}3 & 3\overline{21} & \overline{1}3\overline{2} \end{array} \right\}.$$

Figure 52 depicts these $2+4$ distinct indexed and oriented versions of A and B.



$A(123)$ 24 $A(\overline{1}23)$ 24

$B(123)$ 12 $B(\overline{1}23)$ 12 $B(1\overline{2}3)$ 12 $B(12\overline{3})$ 12

FIGURE 52. The possible entries of a chirotope

Using these notations one can describe the set of chirotopes on the indexing set $\{1,2,3,4\}$ as the set of

$$\chi = \{\chi(123), \chi(124), \chi(134), \chi(234)\}$$

such that, up to a signed permutation of the indices,

(A_1): if $A(213), A(314), A(412) \in \chi$ then $A(234)$ or $A(243)$ or $B(234) \in \chi$;
(A_2): if $B(12\overline{3}), B(12\overline{4}), B(13\overline{4}) \in \chi$ then $B(2\overline{3}4) \in \chi$.

To describe the set of chirotopes on 5 indices we use the coding of an indexed arrangement of oriented pseudolines by its *cycles* : there is exactly one cycle per pseudoline γ of the arrangement and this cycle is defined as the circular sequence of signed indices obtained by writing down the indices of the pseudolines encountered when walking along the side of the pseudoline γ, each index being signed positively or negatively depending on whether the encountered pseudoline is (locally) oriented away or towards the pseudoline γ. For example the cycles of the arrangement $H(12345)$ of Figure 50 are

$$\begin{array}{rl}
1: & 2345\overline{3}\overline{254} \\
2: & 13\overline{4531}45 \\
3: & \overline{1}245\overline{2}1\overline{45} \\
4: & 32\overline{5132}15 \\
5: & 32\overline{1}4\overline{324}1.
\end{array}$$

Similarly the three cycles of $A(123)$ are $23\overline{23}, 31\overline{31}$ and $12\overline{12}$ and those of $B(123)$ are $\overline{2}2\overline{3}3, \overline{3}3\overline{1}1$ and $\overline{1}1\overline{2}2$. The set of chirotopes on a set of 5 indices, say $1,2,3,4,5$, can then be described as the set of

$$\begin{aligned}\chi &= \{\chi(J) : J \in \binom{1,2,3,4,5}{3}\}\\ &= \{\chi(123), \chi(124), \chi(125), \chi(134), \chi(135),\\ &\quad \chi(145), \chi(234), \chi(235), \chi(245), \chi(345)\}\end{aligned}$$

such that the restrictions of χ to the sets of triples of the 5 subsets of 4 indices of $\{1, 2, 3, 4, 5\}$ are chirotopes on 4 indices, i.e., satisfy the axioms (A_1) and (A_2) mentioned above, which satisfy a single additional axiom (A_3) saying that for any index i the 4 cycles indexed by i of these 5 chirotopes on 4 indices are mergeable.

The second and last basic result we need to mention is the following *enlargement theorem for pseudoline arrangements*, due to Goodmann, Pollack, Wenger and Zamfirescu [24], proving a conjecture of B. Grünbaum [27, Conjecture 4.10, page 90].

Theorem 42 ([24])**.** *Any arrangement of pseudolines is an arrangement of lines of a projective plane.* □

Combining the enlargement theorem of pseudoline arrangements with the duality principle for projective planes we get the following theorem.

Theorem 43. *Any arrangement of pseudolines is isomorphic to the dual arrangement of a finite set of points of a projective plane.* □

Proof. Indeed any pseudoline arrangement $\mathcal{A}$ is isomorphic to the dual of the point set $\mathcal{A}$ of the dual projective plane of any projective plane extension of $\mathcal{A}$—here we implicitly use the fact that a projective plane is isomorphic to its bidual. □

APPENDIX B. CHIROTOPES OF FINITE PLANAR FAMILIES OF POINTS

We now review the “classical” characterization of chirotopes of finite planar families of points mentioned in the abstract. (An oriented point in a projective plane is a point together with an orientation of its neighborhood, indicated in ours drawings by an oriented circle surrounding the point.) Our account takes advantage of the relatively recent positive answer of Goodman, Pollack, Wenger and Zamiferescu [24] to the question of Grübaum [27, Conjecture 4.10, page 90] about the embeddability of any arrangement of pseudolines in the line space of a projective plane.

Let Δ be a finite family of oriented points of a projective plane $(\mathcal{P}, \mathcal{L})$, and let τ be a line of $(\mathcal{P}, \mathcal{L})$. We define

(1) the *cocycle of* Δ *at* τ or the *cocycle of the pair* (Δ, τ) as the homeomorphism class of the pair (Δ, τ), i.e., the set of $(\varphi\Delta, \varphi\tau)$ as φ ranges over the set of homeomorphisms of surfaces with domain $\mathcal{P}$; in other terms two pairs (Δ, τ) and (Δ', τ') define the same cocycle if there exists a homeomorphism φ of $\mathcal{P}$ onto $\mathcal{P}'$ such that $\Delta' = \varphi\Delta$ and $\tau' = \varphi\tau$;
(2) the *cocycle-map* as the map that assigns to each cell σ of the dual arrangement of Δ the cocycle of Δ at an element (hence any) of σ;
(3) a 0-, 1, 2-*cocycle* of Δ as a cocycle of Δ at a 0-, 1-, 2-cell of its dual arrangement;
(4) the *isomorphism class of* Δ as the set of configurations Δ' that have the same set of 0-cocycles as Δ (hence,using a simple perturbation argument, the same set of cocycles as Δ); and
(5) the *chirotope of* Δ as the map that assigns to each triple of its indexing set the isomorphism class of the subconfiguration indexed by this triple.

Figure 53 depicts the cocycles of configurations of three points: each circular diagram is labeled at its right bottom corner with its number of reindexations and reorientations and at its left bottom corner with its *cocycle-label* $\nu(\Delta, \tau)$ which is defined as follows. Let $\mathcal{D}_\tau$ be the closed 2-cell obtained by cutting the cross surface $\mathcal{P}$ along the line τ, let $\nu_\tau : \mathcal{D}_\tau \to \mathcal{P}$ be the canonical projection, and let Σ be the pre-image of the set of Δ_is under ν_τ; each element of Σ is endowed with the orientation of the corresponding point of Δ. Choose an orientation ϵ of τ, orient $\mathcal{D}_\tau$ accordingly, and define the *cocycle-label* of the triple (Δ, τ, ϵ) as the set composed of


2
3
1
2
1
3
2
1
3
1
2
3
$132\overline{132}$
4
$12\overline{12}, \overline{3}$
6
$1\overline{1}, \overline{2}, \overline{3}$
6
1, 2, 3
4

FIGURE 53. The cocycles, up to reindexation and reorientation, of configurations of three points with indexing set $\{1, 2, 3\}$

(1) the indices α whose corresponding points Δ_α are contained in the interior of $\mathcal{D}_\tau$ and oriented consistently with the orientation of $\mathcal{D}_\tau$;
(2) the negatives of the indices α whose corresponding points Δ_α are contained in the interior of $\mathcal{D}_\tau$ and oriented inconsistently with the orientation of $\mathcal{D}_\tau$; and
(3) the image of the circular sequence of elements of Σ_τ encountered when walking along the boundary of $\mathcal{D}_\tau$ according to its orientation under the morphism that assigns to a point its index or the negative of its index depending on whether the orientation of the point and the orientation of $\mathcal{D}_\tau$ are consistent or not.

The *cocycle-label of the pair* (Δ, τ) or the *cocycle-label of* Δ *at* τ is then defined as the cocycle-labels of the triples (Δ, τ, ϵ) and $(\Delta, \tau, -\epsilon)$; it can be represented by any of its two elements since the cocycle-label of the triple $(\Delta, \tau, -\epsilon)$ is obtained from the cocycle-label of the triple (Δ, τ, ϵ) by replacing each of its elements by the reversal of its negative (with the usual convention that an index coincides with the negative of its negative : $\overline{\overline{\alpha}} = \alpha$). Clearly the cocycle of a pair (Δ, τ) depends only on its cocycle-label and vice-versa.

A simple case analysis shows that the map that assigns to the isomorphism class of an indexed configuration of oriented points its chirotope is well-defined and one-to-one, that there are exactly six isomorphism classes of indexed configurations of three oriented points on the indexing set $\{1, 2, 3\}$, namely, in cocycle-label terms,

$$\begin{array}{l}
\{123\overline{123}\} \\
\{132\overline{132}\} \\
\{213\overline{213}\} \\
\{1\overline{3}2\overline{1}3\overline{2}\} \\
\{12\overline{12}, 3\}, \{23\overline{2}\overline{3}, 1\}, \{31\overline{3}\overline{1}, 2\} \\
\{12\overline{12}, \overline{3}\}, \{23\overline{23}, \overline{1}\}, \{31\overline{31}, \overline{2}\}
\end{array}$$

and, finally, that the map that assigns to an indexed configuration of three oriented points the isomorphism class of its dual arrangement is compatible with the isomorphism relation on indexed configurations of three oriented points and that the induced (one-to-one and onto) quotient map is the following

$$\begin{array}{lcl}
\{123\overline{123}\} & \longrightarrow & B(1\overline{2}3) \\
\{132\overline{132}\} & \longrightarrow & B(12\overline{3}) \\
\{213\overline{213}\} & \longrightarrow & B(\overline{1}23) \\
\{1\overline{2}3\overline{1}2\overline{3}\} & \longrightarrow & B(123) \\
\{12\overline{12}, 3\}, \{23\overline{2}\overline{3}, 1\}, \{31\overline{31}, 2\} & \longrightarrow & A(123) \\
\{12\overline{12}, \overline{3}\}, \{23\overline{23}, \overline{1}\}, \{31\overline{31}, \overline{2}\} & \longrightarrow & A(\overline{1}23)
\end{array}$$

Since the map that assigns to an isomorphism class of indexed arrangement of oriented pseudolines its chirotope is one-to-one and since any arrangement of pseudolines is isomorphic to the dual arrangement of a family of points it follows that the above considerations concerning indexed configurations of three oriented points and indexed arrangements of three oriented pseudolines extend to configurations of any number of points and arrangements of any number of pseudolines. We summarize

Theorem 44. *The map that assigns to an indexed configuration of oriented points the isomorphism class of its dual arrangement is compatible with the isomorphism relation* *on indexed configurations of oriented points; furthermore the induced quotient map is one-to-one and onto.* □

Therefore there are also six isomorphism classes of cocycle-maps on the indexing set $\{1, 2, 3\}$; they are depicted in Figure 54.



FIGURE 54. The cocycle maps for families of three points indexed by 1, 2 and 3.

The well-informed reader will have recognized here a reformulation, taking advantage of the embeddability of any arrangement of pseudolines in the line space of a projective plane, of the existence of an adjoint—or Type II representation—for every oriented matroid of rank three [22, 10, 23], [2, page 263].

Combining Theorems 43, 44 and 41 we get the characterization of chirotopes of planar families of points mentioned in the abstract.

Theorem 45. *Let χ be a map defined on the set of triples of a finite set I. Then χ is the chirotope of a finite family of oriented points of a projective plane if and only if for every 3-, 4-, and 5-subset J of I the restriction of χ to the set of triples of J is the chirotope of a finite family of pairwise disjoint oriented points of a projective plane.* □

Appendix C. Convexity in neutral and projective planes

In this section we prove that the polar of a convex body is a convex body, that the dual of a convex body is the boundary of its polar (hence a double pseudoline) and that, up to homeomorphism, the dual arrangement of a pair of disjoint convex bodies is the unique arrangement of two double pseudolines that intersect in four transversal intersection points and induce a cellular decomposition of their underlying cross surface.

We proceed in two steps, establishing the properties first for (real two-dimensional) neutral planes, which we shall briefly recall, and second for projective planes by reduction to the first step.

C.1. **Background material on neutral and affine planes.** A (real two-dimensional) *neutral plane* is a topological point-line incidence geometry $(\mathbb{A}, \mathbb{L})$ whose point space $\mathbb{A}$ is homeomorphic to $\mathbb{R}^2$, whose line space $\mathbb{L}$ is a subspace of the space of pseudolines of $\mathbb{A}$, and whose axiom system is reduced to the following single axiom: any two distinct points belong to exactly one line, called their *joining line*, which depends continuously on the two points.[2]

The *join map*, denoted $\vee$, assigns to any ordered pair of distinct points of $\mathbb{A}$ their joining line in $\mathbb{L}$; the *intersection map*, denoted $\wedge$, assigns to any ordered pair of distinct intersecting lines of $\mathbb{L}$ their common intersection point in $\mathbb{A}$. The join and intersection maps are continuous and open.

Theorem 46 ([6, page 220]). *Let $\mathbb{L}$ be the line space of a neutral plane and let $\widetilde{\mathbb{L}}$ be the space of oriented versions of the lines of $\mathbb{L}$. Then $\mathbb{L}$ is an open Möbius strip, the natural projection $\widetilde{\mathbb{L}} \to \mathbb{L}$ that assigns to an oriented line is unoriented version is a two-covering map, and the pencil of lines through a point is a pseudoline in $\mathbb{L}$, i.e., a nonseparating simple closed curve embedded in $\mathbb{L}$.* □

An *affine* plane is a neutral plane with the property that for every point-line pair (p, ℓ) there exists a unique line k incident to p such that either $k = \ell$ or k and ℓ are disjoint; the line k is called the *parallel* to ℓ through p and the set of parallels to ℓ, denoted $[\ell]$, is called the *parallel class* of ℓ. Note that the set of parallel classes is a partition of the line space of the plane. The *projective completion* of an affine plane $(\mathbb{A}, \mathbb{L})$ is the projective plane whose line space $\widehat{\mathbb{L}}$ and point space $\widehat{\mathbb{A}}$ are, respectively,

(1) the set $\{\ell \cup \{[\ell]\} \mid \ell \in \mathbb{L}\} \cup \{[\mathbb{L}]\}$, where $[\mathbb{L}]$ is the set of parallel classes, endowed with the topology of the one-point compactification $\mathbb{L} \cup \{\infty\}$ of $\mathbb{L}$ via the map that assigns $\ell \in \mathbb{L}$ with $\ell \cup [\ell] \in \widehat{\mathbb{L}}$ and ∞ with $\{\mathbb{L}\}$; and
(2) the set $\mathbb{A} \cup [\mathbb{L}]$ endowed with the topology with subbase the $J_p \wedge J_q$ where p and q are two points of $\widehat{\mathbb{A}}$ and where J_p and J_q are disjoint open intervals of the pencils of lines through p and q, respectively.

Theorem 47. *The projective completion of an affine plane is a projective plane.* □

---

[2] The space of pseudolines of $\mathbb{A}$ is the quotient of the space of embeddings of $\mathbb{R}$ into $\mathbb{A}$ with closed images (i.e., the set of continuous one-to-one maps $\varphi : \mathbb{R} \to \mathbb{A}$, with the property that $\varphi(\mathbb{R})$ is closed in $\mathbb{A}$, endowed with the compact-open topology) under the natural action of the group of homeomorphisms of $\mathbb{R}$. As usual we identify a pseudoline φ with its image $\varphi(\mathbb{R})$. Similarly the space of oriented pseudolines of $\mathbb{A}$ is the quotient of the space of embeddings of $\mathbb{R}$ into $\mathbb{A}$ with closed images under the natural action of the group of direct homeomorphisms of $\mathbb{R}$.

The *affine part* of a real two-dimensional projective plane $(\mathcal{P}, \mathcal{L})$ at $\ell \in \mathcal{L}$ is the topological point-line incidence geometry $(\mathcal{P} \setminus \ell, \mathcal{L} \setminus \{\ell\})$.

Theorem 48. *The affine parts of a projective plane are affine planes with the property that their projective completions are isomorphic to the initial projective plane.* □

We refer to the monograph of Salzmann et al [40, Chap. 3] for supplementary background material on neutral planes, where they are called $\mathbb{R}^2$-planes.

C.2. **Convexity and duality in (real two-dimensional) neutral planes.** We work in a neutral plane $(\mathbb{A}, \mathbb{L})$. As in the Euclidean plane, a subset of points is called *convex* if it includes the line-segments joining its points. A compact convex subset of points with nonempty interior is called a *convex body*, and a *tangent* or a *supporting line* to a convex body is a line that intersects the convex body but not its interior (equivalently, a line that intersects the convex body and that includes the body in one of its two closed sides). Finally we define the dual U^* of a convex body U as its set of tangent lines.

In this section we establish the following basic twofold result.

Theorem 49. *The dual of a convex body of a* $\mathbb{R}^2$*-plane* $(\mathbb{A}, \mathbb{L})$ *is a double pseudoline in* $\mathbb{L}$ *(i.e., a nontrivial separating simple closed curve embedded in* $\mathbb{L}$*) and the dual family of a family of two disjoint convex bodies is an arrangement of two double pseudolines that intersect in four transversal intersection points and induce a cellular decomposition of (the one-point compactification of)* $\mathbb{L}$*.* □

The proof proceeds by a sequence of auxiliary results.

C.2.1. *Boundary of a convex body and tangents.* The proofs of the two following lemmas are adapted from [1, Chap. 11.3].

Lemma 50. *The boundary of a convex body is a simple closed curve.* □

Proof. Let U be a convex body, let A be one of its interior point, and let $L_A \approx \mathbb{S}^1$ be the pencil of oriented lines through A. Consider the application $\varphi : L_A \to \partial U$ that assigns to $\ell \in L_A$ the endpoint of the trace of U on ℓ beyond A. Clearly φ is a well-defined one-to-one and onto correspondence whose inverse is continuous. Therefore it is sufficient to show that φ is continuous. Let $\ell \in L_A$ and let B and C be two points of the interior of U with A contained in the interior of the line segment joining B to C. As illustrated in Figure 55 the half-lines with origins A, B and C through $\varphi(\ell)$ leave U at $\varphi(\ell)$. Let $\{\ell_n\}$, $n \in \mathbb{N}$, be a sequence of oriented lines of L_A converging to ℓ with $\ell_n \neq \ell$. For n large enough the intersection points of the line ℓ_n with the lines $B \vee \varphi(\ell)$ and $C \vee \varphi(\ell)$ are well-defined and converge to $\varphi(\ell)$. Consequently, for n large enough, the line ℓ_n intersects the half-lines through $\varphi(\ell)$ with origins B and C. One of these two intersection points is beyond $\varphi(\ell)$ and the other one is before $\varphi(\ell)$. Therefore $\varphi(\ell_n)$ belongs to the line segment joining these two points. When n goes to infinity, this line segment retracts onto $\varphi(\ell)$. Therefore φ is continuous. □

Lemma 51. *Let* U *be a convex body and let* A *be a point not in the interior of* U*. Then the set* S *of lines through* A *intersecting the interior of* U *is an open interval of the pencil of lines through* A*. Furthermore*



FIGURE 55.

(1) *if $A \notin \partial U$, then the set T of lines through A missing U is an open interval of the pencil of lines through A whose endpoints are the endpoints of S, and the set of tangents to U through A is a pair composed of the two common endpoints of S and T;*

(2) *if $A \in \partial U$, then the set of tangents to U through A is a closed interval of the pencil of lines through A whose endpoints are the endpoints of S.* □

Proof. The set S is

(1) an open subset of the pencil of lines through A, because the join map is open;
(2) connected because the interior of U is connected.

Consequently S is an open interval of the pencil of lines through A or S is the whole pencil of lines through A.

We now show that S is not the pencil of lines through A.

For $\ell \in S$ let ℓ^+ and ℓ^- be the two connected components of $\ell \setminus \{A\}$ with the convention that ℓ^+ is the connected component that intersects the interior of U (and, consequently, ℓ^- is the connected component that does not intersect U). We set $R^+ = \bigcup S^+$ and $R^- = \bigcup S^-$.

Let $\mathcal{I}$ be the set of closed line segments I contained in the interior of U whose supporting line is not incident to A and for any $I \in \mathcal{I}$ let S_I be the set of lines of S intersecting the interior of I. We set $Q^+(I) = \bigcup S_I^+$, and $Q^-(I) = \bigcup S_I^-$. We claim that

(1) $\mathcal{I}$ is nonempty;
(2) $S = \bigcup_{I \in \mathcal{I}} S_I$;
(3) $Q^+(I)$ and $Q^-(I)$ are open quarters of plane;
(4) $R^+ = \bigcup_{I \in \mathcal{I}} Q^+(I)$ is open and nonempty;
(5) $R^- = \bigcup_{I \in \mathcal{I}} Q^-(I)$ is open and nonempty.

The sets R^+ and R^- are disjoint nonempty open subsets of the plane minus the point A. This last set being connected, there exists a point E not in R^+ nor in R^-. The line joining A and E misses the interior of U. Consequently S is not the pencil of lines through A.

The endpoints of the interval S intersect the boundary of U because U is compact, and U is contained in only one of the two sides delimited by these endpoints (otherwise we can find a triangle contained in U with two of its vertices separated by the line, and the line will intersect the interior of U). ☐



Figure 56.

C.2.2. *Duality.*

Lemma 52. *Let U be a convex body and let U^* be its set of tangents. Then*

1. *U^* is a simple closed curve;*
2. *U^* is a double pseudoline in $\mathbb{L}$;*
3. *the set of lines missing U is the one-punctured topological disk bounded by U^*;*
4. *the set of lines intersecting the interior of U is the Möbius strip bounded by U^*.*

Proof. Let O be an orientation of the plane, let Δ be the map that assigns to A not in U the tangent to U through A with the property that walking along the tangent from A to U we see the convex body U on our right, let $I = [A, B]$ be a closed line segment missing U with the property that $\Delta(A) \neq \Delta(B)$, and let Γ be a simple closed curve surrounding U. We claim that

1. Δ is continuous and onto;
2. the restriction of Δ to the domain Γ is onto;
3. the restriction of Δ to the domain I and codomain $\Delta(I)$ is a homeomorphism;
4. Δ is open;

from which it follows that U^* is a compact 1-dimensional manifold, hence a simple closed curve.

We now prove Claims (2), (3), and (4). According to Claim (1) the set U^* is a simple closed curve of $\mathbb{L}$ and, consequently, it is a pseudoline or a double pseudoline of the one-point compactification $\mathbb{L} \cup \{\infty\}$ of $\mathbb{L}$. Now recall that two pseudolines intersect necessarily in at least one point and observe that the pencil of lines through a point lying in the interior of U is a pseudoline that does not intersect U^*, to conclude that U^* is a double pseudoline of $\mathbb{L} \cup \{\infty\}$. Finally the set of lines intersecting the interior of U is a connected component of the complement of U^* that contains pseudolines. Consequently it is a Möbius strip in $\mathbb{L} \cup \{\infty\}$. Now U being compact the set of lines intersecting U is compact in $\mathbb{L}$ (cf. [40, page 160]). Consequently the Möbius strip bounded by U^* does not contain ∞. This proves claims (2), (3) and (4). ☐

Lemma 53. *Let U and V be two disjoint convex bodies. Then the double pseudolines U^* and V^* intersect in exactly four points, where they cross.* ☐

Proof. Let O be an orientation of the plane, let Δ be the map that assigns to $A \notin U$ the tangent to U through A with the property that walking along the tangent from A to U we see the convex body U on our right, let $I = [A, B]$ be a closed line segment missing U with the property that $\Delta(A) \neq \Delta(B)$, and let Γ be a simple closed curve surrounding U. We claim that

(1) Δ is continuous and onto;
(2) the restriction of Δ to the domain Γ is onto;
(3) the restriction of Δ to the domain I and codomain $\Delta(I)$ is a homeomorphism;
(4) $\Delta(V)$ is a closed interval $[T, T']$, $T \neq T'$, of U^*;
(5) $\Delta(\mathrm{Int}(V))$ is the open interval $]T, T'[$;

from which it follows that T and T' are the sole tangents to both U and V such that walking along the tangents from V to U we see U on our right (and walking along the tangents from U to V we see V on our left or on our right depending on whether we walk on T or on T'), and that U^* and V^* cross at T and T'. Permuting the role of U and V we get a second pair of common tangents to U and V. This proves the lemma. ☐

Lemma 54. *Let U and V be two disjoint convex bodies. Then the double pseudolines U^*, V^* induce a cellular decomposition of the (one-point compactification of) $\mathbb{L}$.* ☐

Proof. Let $u \in \mathrm{Int}(U)$ and let $v \in \mathrm{Int}(V)$. We have seen that

(1) U^* and V^* are double pseudolines intersecting in exactly four points—where they cross;
(2) u^* and v^* are pseudolines intersecting in exactly one point—where they cross;
(3) u^* is contained in the Möbius strip bounded by U^*;
(4) u^* and V^* intersect in exactly two points—where they cross;
(5) v^* is contained in the Möbius strip bounded by V^*;
(6) v^* and U^* intersect in exactly two points—where they cross.

Consequently the arrangements $\{u^*, v^*, U^*\}$ and $\{u^*, v^*, V^*\}$ are (up to homeomorphism) the ones depicted in the two first diagrams of Figure 57. Now it is not hard to see that the only compatible superpositions of these two diagrams, fullfilling condition (1) above, are the two last diagrams of Figure 57. It remains to observe that the set of lines intersecting both U and V is connected to rule out the last diagram from our considerations and to conclude that the pair $\{U^*, V^*\}$ is an arrangement of double pseudolines. ☐



Figure 57.

Observe that this proves that two disjoint convex bodies have a strictly separating line.

C.3. **Convexity and duality in projective planes.** We now work in a projective plane $\mathcal{G} = (\mathcal{P}, \mathcal{L})$.

Lemma 55. *Assume that two of the three sides of a triangular face of a simple arrangement of three lines are contained in a convex body. Then the triangular face is contained in the convex body.* $\square$

Proof. Let O be a compact subset of points with nonempty interior whose intersection with any line is empty, a line segment or a full line and let T be a triangular face of a simple arrangement of three lines. Let A, B, C be the three vertices of the triangular face T, as illustrated in the left part of Figure 58 where the triangular face is marked with a little square, let $[AB], [BC]$ and $[CA]$ be the three sides of T and assume that $[AB]$ and $[AC]$ are contained in O. Proving our lemma comes down to proving that O contains a line or that T is contained in O. Let D be a point of the line (BC) outside the line segment $[BC]$, as illustrated in the right part of Figure 58. For any $X \in [AB]$, we denote by X' the intersection of the line (DX) with the line (AC)—note that X' ranges over the line segment $[AC]$ and that $B' = C$—by $[XX']$ the line segment supported by the line (DX) contained into T, and, for $X \neq A$, by $[X'X]$ the line segment of (DX) contained in the complement of the interior of T. Let I_1 be the set of $X \in [AB]$, $X \neq A$, such that $[XX']$ is contained in O, and let I_2 be the set of $X \in [AB]$, $X \neq A$, such that $[X'X]$ is contained in O. One can easily check that



Figure 58.

(1) $I_1 \cap I_2 = \emptyset$ unless O contains a line (DX) with $X \in [AB], X \neq A$;
(2) $I_1 \cup I_2 = [AB] \setminus \{A\}$;
(3) I_1 and I_2 are both closed in $[AB] \setminus \{A\}$.

Assume now that $I_1 \cap I_2 = \emptyset$, otherwise O contains a line (DX) with $X \in [AB], X \neq A$, and we are done. Since $[AB] \setminus \{A\}$ is connected, it follows that I_2 or I_1 is empty. In the first case $T = \bigcup_{X \in [AB]} [XX'] \subseteq O$ and in the second case we see easily, since O is compact, that the line (DA) is contained in O. In both case we are done. $\square$

Lemma 56. *The trace of a line on the interior of a convex body is empty or is the interior of the trace of the line on the body.* $\square$

Proof. Let O be a convex body, let $[AB]$ be the trace of a line on O and assume that $[AB]$ intersects the interior of O at point C. Let $[DE]$ be a line segment through C, contained in the interior of O, and not contained in the line (AB). Let $[AC]$ be the line segment with endpoints A and C contained in $[AB]$. Similarly let $[BC]$ be the line segment with endpoints B and C contained in $[AB]$. In view of Lemma 55 the four triangles with sides $[AC]$ and $[CD]$, $[AC]$ and $[CE]$, $[BC]$ and $[CD]$, $[BC]$ and $[CE]$ are contained in O. See Figure 59 for an illustration.



FIGURE 59.

$\square$

Let ℓ be a line of $(\mathcal{P}, \mathcal{L})$, let $\mathcal{D}$ be the closed topological disk obtained by cutting $\mathcal{P}$ along ℓ, and let $\mathcal{D} \to \mathcal{P}$ be the induced canonical map. Let O be a convex body and let $\widetilde{O}$ be the pre-image of O under $\mathcal{D} \to \mathcal{P}$. If the line ℓ misses O then O is a convex body of the affine part of $(\mathcal{P}, \mathcal{L})$ obtained by removing the line ℓ and $O = \widetilde{O}$. We now examine the situation when ℓ intersects O. Let I be the trace of ℓ on O, and assume that I is nonempty. We denote by I^+ and I^- the two connected components of the pre-image of I under $\mathcal{D} \to \mathcal{P}$. See Figure 60 for an illustration. For all $A, B \in O$, $A \neq B$, we denote by $[A, B]_O$ the line segment joining A and B contained in O. Let $A \in I$. For all $B \in O \setminus I$ the pre-image under $\mathcal{D} \to \mathcal{P}$ of $[A, B]_O$ has two connected components: a point A_B^* and $([A, B]_O \setminus \{A\}) \cup A_B$ where $\{A_B^*, A_B\}$ is the pre-image of A under $\mathcal{D} \to \mathcal{P}$. Let O^+ be the set of $B \in O \setminus I$ such that $A_B \in I^+$ and, similarly, let O^- be the set of $B \in O \setminus I$ such that $A_B \in I^-$. Clearly by definition $O^+ \cup O^- = O \setminus I$ and $O^+ \cap O^- = \emptyset$.



FIGURE 60.

Lemma 57. *We claim that*

1. *O^+ and O^- are independent of the choice of $A \in I$;*
2. *O^+ and O^- are closed convex subsets of the affine part of $(\mathcal{P}, \mathcal{L})$ obtained by removing the line ℓ;*
3. *O^+ and O^- are both nonempty if and only if I intersects the interior of O;*
4. *the topological closure of O^+ in $\mathcal{D}$ is $O^+ \cup I^+$ under the assumption that O^+ is nonempty, and a similar result holds for O^-.*

In particular $\widetilde{O}$ has two connected components: $O^+ \cup I^+$ and $O^- \cup I^-$. □

Proof. Claims (1), (2) and (3) are simple applications of Lemmas 55 and 56. Assume now that O^+ is nonempty. Clearly the topological closure of O^+ contains $O^+ \cup I^+$ and is contained in $O^+ \cup I^+ \cup I^-$. Thus we have to prove that the topological closure of $O^+ \cup I^+$ avoids I^-. Assume the contrary. Then there exists a convergent sequence of points $A_n \in O^+$ with limit $A' \in I^-$. Let $A \in I$, let $J_n = [A, A_n]_O \setminus \{A\}$ and let ℓ_n be the supporting line of J_n. Without loss of generality one can assume that the sequence ℓ_n has a limit ℓ'. Let $B \in \ell'$, $B \notin I$. There exists a convergent sequence of points $B_n \in \ell_n$ with limit B. For n large enough B_n belongs to J_n. Since O^+ is compact it follows that $B \in O^+$ and consequently ℓ' is a subset of O. A contradiction with the assumption that O is a convex body. □

Lemma 58. *Assume that there a line missing the interior of a convex body. Then there is a line missing the body.* □

Proof. Let O be a convex body, let ℓ be a line missing the interior of O, let I be the trace of ℓ on O and assume that I is nonempty (otherwise we are done). Let ℓ_∞ be a line missing the line segment I and let Q and Q' be the two connected components of the complement of the lines ℓ and ℓ_∞ in $\mathcal{P}$, as indicated in the left part of Figure 61. Let U be a neighboorhood of the intersection point of ℓ and ℓ_∞ disjoint from O. Let V_n be a decreasing sequence of open neighborhoods of ℓ with $\bigcap_n V_n = \ell$ and let W_n and W'_n be the traces of V_n on $Q \cup U$ and $Q' \cup U$, respectively. According to the previous lemma there exists an n_0 such that for all $n \geq n_0$ the trace of W_n on O is empty or for all $n \geq n_0$ the trace of W'_n on O is empty. Without loss of generality one can assume that the trace of W'_n on O is empty for $n \geq n_0$. Using standard compactness arguments we see that there is a line ℓ'' of the pencil of line through the intersection point of ℓ and ℓ_∞ contained in W'_n and we are done. □



Figure 61.

Lemma 59. *Assume that there a line missing the interiors of two disjoint convex bodies. Then there is a line missing the two bodies.* ☐

Proof. Let O and O' be two disjoint convex bodies, let ℓ be a line missing the interiors of O and O', let I and J be the traces of ℓ on O and O' and assume that I and J are nonempty (otherwise we are done, thanks to the previous lemma), as indicated in the left part of Figure 62. Let $\mathcal{D}$ be the closed topological disk obtained by cutting $\mathcal{P}$ along ℓ, let $\mathcal{D} \to \mathcal{P}$ be the induced canonical map, let I^+ and I^- be the two connected components of the pre-image of I under $\mathcal{D} \to \mathcal{P}$ with the convention that I^- is also a connected component of the pre-image of O under $\mathcal{D} \to \mathcal{P}$, and similarly let J^+ and J^- be the two connected components of the pre-image of J under $\mathcal{D} \to \mathcal{P}$ with the convention that J^- is also a connected component of the pre-image of O under $\mathcal{D} \to \mathcal{P}$. The lemma follows the simple observation that there is a line misssing I and J whose pre-image under $\mathcal{D} \to \mathcal{P}$ separates $I^+ \cup J^+$ from $I^- \cup J^-$, as indicated in the right part of Figure 62. ☐



FIGURE 62.

Lemma 60. *Any boundary point of a convex body is incident to a line missing the interior of the body.* ☐

Proof. Let O be a convex body and let A be a boundary point of O. The color of an oriented line ℓ incident to the boundary point A is defined to be

1. *blue* if the line ℓ does not intersect the interior of O;
2. *white* if the line ℓ intersects the interior of O and if A is the initial point of the trace on the interior of O of the oriented line ℓ;
3. *red* if the line intersects the interior of O and if A is the terminal point of the trace on the interior of O of the oriented line ℓ.

According to Lemma 56 any oriented line through A has a color, and these colors are mutually exclusive. The sets of white and red oriented lines are open subsets of the pencil of oriented lines through A. Since none of these two sets is empty and since the pencil of oriented lines through A is connected it follows that the set of blue oriented lines is nonempty. This proves the lemma. ☐

Combining Lemmas 58 and 60 we get that the set of lines missing a convex body is nonempty.

Lemma 61. *The set of lines missing a convex body is nonempty.* ☐

Let O be a convex body of $\mathcal{G}$ and let ℓ be a line missing O. Working in the affine part $\mathcal{G}^\ell$ of $\mathcal{G}$ we see that a convex body of a projective plane is also a convex body of an affine $\mathbb{R}^2$-plane. In particular there exists a tangent through any boundary point and the set of tangents through a boundary point is a segment of the pencil of lines through that boundary point.

Lemma 62. *The set of lines missing two disjoint convex bodies is nonempty.* ☐

Proof. According to Lemma 59 it is sufficient to prove that the set of lines missing the interiors of two disjoint convex bodies is nonempty. Let U and V be two disjoint convex bodies and let ℓ be a line missing U. If ℓ avoids the interior of V we are done. Otherwise ℓ intersects V along a closed line segment, say $[RS]$, $R \neq S$, and ℓ intersects the interior of V along the interior of $[RS]$.



Figure 63.

Let τ be a tangent to V at S, let ν be a tangent to V at R and let A be the intersection point of τ and ν. If τ or ν misses the interior of U we are done. Otherwise we proceed as follows.

Let g be a line through A that avoids the interior of V, let B be the intersection point of ℓ and g, let $h \neq \ell$ be a line through B that avoids U but intersects the interior of V, and let W be the intersection of V with the strip delimited by g and h in the affine geometry $(\mathcal{P} \setminus \ell, \mathcal{L} \setminus \{\ell\})$. Clearly U and W are disjoint convex bodies of the affine geometry $(\mathcal{P} \setminus \ell, \mathcal{L} \setminus \{\ell\})$: Let D be the intersection point of their interior bitangents. We let the reader check that D belongs to the triangle in $\mathcal{P} \setminus \ell$ delimited by the lines τ, ν and h and that the line through D of the pencil of lines through B avoids the interiors of U and V. ☐

Luc Habert
E-mail address: `Luc.Habert@normalesup.org`

Michel Pocchiola, Université Pierre & Marie Curie, Institut de Mathématiques de Jussieu (UMR 7586), 4 place Jussieu, 75252 Paris Cedex, France
E-mail address: `pocchiola@math.jussieu.fr`